arXiv:1205.2905v3 [math.DS] 19 Nov 2012

UNIQUENESS OF THE MEASURE OF MAXIMAL ENTROPY FOR THE SQUAREFREE FLOW

RYAN PECKNER

Abstract. The squarefree flow is a natural dynamical system whose topological and ergodic properties are closely linked to the behavior of squarefree numbers. We prove that the squarefree flow carries a unique measure of maximal entropy and describe the structure of the associated measure-preserving dynamical system. Our method involves first studying approximations arising from finite collections of prime numbers, then taking a limit under Ornstein's $\bar{d}$-metric in order to consider all primes simultaneously. This is accomplished by proving uniform Gibbs bounds for a sequence of sofic systems and constructing explicit joinings between them in order to directly estimate their $\bar{d}$-distances.

1. Introduction and Background

One of the most important objects in analytic number theory is the Möbius function, defined for positive integers n by

$$\mu(n) = \begin{cases} 1 & \text{if } n = 1 \\ 0 & \text{if } n \text{ is not squarefree} \\ (-1)^r & \text{if } n = p_1 \cdots p_r. \end{cases}$$

It is well-known that if the sequence $(\mu(n))_{n\geq 1}$ behaves randomly in the sense that

$$\sum_{n\leq N} \mu(n) = O_\epsilon\left(N^{1/2+\epsilon}\right) \quad \text{for any } \epsilon > 0,$$

then the Riemann hypothesis is true. Less quantitative but richer reflections of the chaotic behavior of μ are captured by instances of the "Möbius Randomness Law", see for example [IK].

Sarnak has recently developed an approach to the idea of Möbius randomness based on the study of certain dynamical systems [Sar]. Let $\Omega^{(3)} = \{-1, 0, 1\}^{\mathbb{N}}$ with the product topology, and let $T : \Omega^{(3)} \to \Omega^{(3)}$ be the left shift defined by $(Tx)_n = x_{n+1}$. $(\Omega^{(3)}, T)$ is a topological dynamical system, that is, a pair (X, S) where X is a compact metric space and $S : X \to X$ is continuous and surjective.

The Möbius function defines a point $\omega = (\mu(1), \mu(2), \mu(3), \dots) \in \Omega^{(3)}$, and we let $\mathcal{M}$ be the closure in $\Omega^{(3)}$ of the orbit $\{T^k\omega : k \geq 0\}$. Then the 'Möbius flow' $\mathcal{M}$ is a closed, T-invariant subset of $\Omega^{(3)}$, but due to the mysterious behavior of μ one cannot say much else about it. The purpose of this paper is to study a related system about which we can say a great deal.

Let $\Omega^{(2)} = \{0, 1\}^{\mathbb{N}}$ and let $\phi : \Omega^{(3)} \to \Omega^{(2)}$ be the squaring map $(x_n) \mapsto \left(x_n^2\right)$. Observe that ϕ is continuous, surjective, and intertwines the shift maps on the two spaces (this is an example of a *factor map* between topological dynamical systems). Let $\mathcal{S}$ be the closure in $\Omega^{(2)}$ of the orbit of $\phi(\omega) = \left(\mu^2(n)\right)_{n\geq 1}$ under T. Since $\mu^2(n)$ is the indicator function of the set of squarefree numbers, we call $(\mathcal{S}, T)$ the squarefree flow.

Naturally, the dynamical structure of $\mathcal{S}$ is strongly tied to the statistical properties of squarefree numbers. In this connection, Sarnak uses the well-known squarefree sieve (cf. [Tsa]) to produce a certain ergodic measure ν of zero entropy on $\Omega^{(2)}$ whose support is $\mathcal{S}$ (for the definitions of terms from ergodic theory see [Gla]). This allows him in turn to prove the following: define the *support* of a sequence $x = (x_n) \in \Omega^{(2)}$ to be the set $\operatorname{supp}(x) \subset \mathbb{N}$ of those n such that $x_n = 1$. We call a subset $A \subset \mathbb{N}$ *admissible* if for every prime p the reduction $A \bmod p^2$ is a proper subset of $\mathbb{Z}/p^2\mathbb{Z}$. We then call a sequence $(x_n) \in \Omega^{(2)}$ admissible if its support is an admissible subset of $\mathbb{N}$. It should be clear that $(\mu^2(n))_{n\geq 1}$ is admissible in this sense.

Let $\mathcal{A} \subset \Omega^{(2)}$ be the set of all admissible sequences. It's easy to show that $\mathcal{A}$ is a closed, shift invariant subset of $\Omega^{(2)}$. Using the aforementioned measure ν (and in particular the fact that its support is $\mathcal{S}$), Sarnak is able to show that in fact $\mathcal{A} = \mathcal{S}$. Though this may seem surprising at first, it arises quite naturally from the shape of the measure ν, which in turn is constructed from the main term of the squarefree sieve formula.

The ergodicity of ν implies that the subset of $\mathcal{A}$ consisting of sequences whose support omits only one residue class $\bmod p^2$ for every p has full ν-measure. It's then not hard to show that the map which sends such a sequence to the point in $\mathcal{K} := \prod_p \mathbb{Z}/p^2\mathbb{Z}$ whose p-th coordinate is the residue class omitted by the sequence $\bmod p^2$ intertwines the shift on $\mathcal{A}$ with the translation on $\mathcal{K}$ by the element $(-1, -1, -1, \dots)$ (note that this map is Borel measurable but not continuous). Moreover, the pushforward of ν under this map is precisely the mass one Haar measure on the compact group $\mathcal{K}$, so we have a factor map of *measure-preserving* dynamical systems $(\mathcal{A}, T, \nu) \to (\mathcal{K}, T_{(-1,-1,-1,\dots)}, \nu_{\text{Haar}})$. Subsequently, Cellarosi and Sinai [CS] used spectral theory techniques to prove that this map is in fact an isomorphism of measure-preserving dynamical systems (it is however very far from being an isomorphism of *topological* dynamical systems).

While this tells the whole story for the measure ν, there are other invariant measures on $\mathcal{A}$ that should be taken into account. Specifically, as stated by Sarnak (and proven in this paper), the system $(\mathcal{A}, T)$ has topological entropy $(6/\pi^2)\log 2$ (it's no coincidence that $6/\pi^2$ is the density of the squarefree numbers in $\mathbb{N}$!). The variational principle ([Gla] Thm. 17.1) states that the topological entropy of a topological dynamical system is the supremum of the measure entropies over all its invariant probability measures, and it is known that any subshift system possesses at least one *measure of maximal entropy*, i.e. an invariant probability measure whose measure entropy equals the topological entropy of the system. Thus there exists such a measure for $\mathcal{A}$.

While measures of maximal entropy always exist for subshifts (this is not the case for all topological dynamical systems), the number of such measures may be arbitrarily large, even infinite (see [Hay] for examples). In keeping with the general principle in ergodic theory that a scarcity of measures is more meaningful than an abundance of them, it is an important problem to determine when there is only one measure of maximal entropy on a given topological dynamical system. This uniqueness can be of great practical value. For instance, the existence of a unique measure of maximal entropy for the geodesic flow on the modular surface is a key ingredient in the proof of Linnik's theorem on the representations of integers by integral binary quadratic forms [Ei].

The aim of this paper is to prove the following.

Theorem (Thm. 3.10). *The squarefree flow $\mathcal{A}$ possesses a unique measure of maximal entropy.*

Our proof relies on a detailed study of the finite-level systems

$$\mathcal{A}(r) := \{x \in \Omega^{(2)} : \operatorname{supp}(x) \bmod p_i^2 \text{ is a proper subset of } \mathbb{Z}/p_i^2\mathbb{Z} \text{ for } i = 1, \ldots, r\}$$

where $p_1, \ldots, p_r$ are the first r primes. Our first result (Prop. 2.5 below) is that each of these systems is a so-called *sofic* shift, meaning they satisfy a certain finitary condition that controls the complexity of the words appearing in the shift (see section 2.2). Such shifts were first investigated by Weiss ([Wei]), who proved that a subshift is sofic if and only if it is a factor of a subshift of finite type (none of the $\mathcal{A}(r)$ is of finite type). It should be mentioned that although Weiss was the first to give an explicit definition, versions of sofic systems were known perhaps long before him to researchers in theoretical computer science; indeed, sofic systems can also be characterized as essentially those subshifts that are recognizable by a finite automaton (see the excellent book [LM] for this perspective).

Sofic shifts are among the few classes of systems that are known to carry a unique measure of maximal entropy. However, while the shifts $\mathcal{A}(r)$ are all sofic the full squarefree system $\mathcal{A} = \cap_{r=1}^{\infty} \mathcal{A}(r)$ is not, so we have to take a different approach to proving that it has a unique maximal measure. Naturally, we do this by exhibiting $\mathcal{A}$ as a certain limit of the systems $\mathcal{A}(r)$ with their measures of maximal entropy, specifically as their limit under Ornstein's $\bar{d}$ metric (see section 3.3 for precise definitions). Philosophically, this metric judges the closeness of two measures on a given shift space by asking how much on average one has to change a generic sequence for the first measure in order to get a generic sequence for the second measure. Limits in $\bar{d}$ are much more complex than weak* limits. In particular, the ergodic measures in the set of invariant probability measures on a shift space are weak* dense but closed in $\bar{d}$, and this is one of the key facts we use to establish the intrinsic ergodicity of $\mathcal{A}$.

Our proof that the systems $\mathcal{A}(r)$ with their respective unique measures of maximal entropy η_r converge under $\bar{d}$ to $(\mathcal{A}, \eta)$ (where η is the weak* limit of the η_r) proceeds by explicitly constructing certain joinings between these systems that reflect their common algebraic origin. More precisely, we show the following in section 3.2.

Proposition (Prop. 3.9)**.** *For each $r \geq 1$, there exists a natural factor map $\phi_r : \left(\mathbb{Z}_r \times \Omega^{(2)}, m_r \times B(\frac{1}{2}, \frac{1}{2})\right) \to (\mathcal{A}(r), \eta_r)$.*

In the above, m_r denotes the counting measure on $\mathbb{Z}_r$ (equipped with translation by $(-1, -1, \ldots, -1)$) and $B(\frac{1}{2}, \frac{1}{2})$ is the maximum entropy Bernoulli measure on $\Omega^{(2)}$. We therefore get the joining $\rho_{rs} := (\pi_{rs} \circ \phi_r \times \phi_s)_* \left(m_s \times B(\frac{1}{2}, \frac{1}{2})\right)$ of $(\mathcal{A}(r), \eta_r)$ and $(\mathcal{A}(s), \eta_s)$ (where $\pi_{rs} : \mathbb{Z}_s \times \Omega^{(2)} \to \mathbb{Z}_r \times \Omega^{(2)}$ is the natural projection), which allows us to show that the $\bar{d}$-distance between them is small. In fact it yields the following bound for the $\bar{d}$-distance, which reflects the algebraic origin of the systems in terms of squares of the prime numbers $p_1 < p_2 < \cdots$.

Proposition (§3.2)**.** *For any $s > r$ we have*

$$\bar{d}(\eta_r, \eta_s) \leq \frac{1}{2}\left(\frac{1}{p_{r+1}^2} + \frac{1}{p_{r+2}^2} + \cdots + \frac{1}{p_s^2}\right).$$

This shows that $\bar{d}(\eta_r, \eta_s) \to 0$ as $r, s \to \infty$.

While these constructions are quite specific to our setting, they reflect a more general principle that certain properties (in this case, intrinsic ergodicity) are preserved under taking the limit of a rapidly converging (in a suitable sense) decreasing sequence of systems with that property. Similar ideas are explored from a very different direction in [Gur]. In this vein, we prove a uniformity in the so-called lower Gibbs bounds for the measures η_r, which effectively states that, up to a constant, the η_r-measure of a cylinder set defined by a word in $\mathcal{A}(r)$ is determined by the word's length. This is a much stronger form of the asymptotic equipartition property guaranteed for any ergodic measure by the Shannon-McMillan-Breiman theorem (cf. [Gla]§14.5). Specifically, we show the following by closely analyzing the definition of the measure of maximal entropy for a sofic system in terms of the Perron-Frobenius eigenvector of a certain adjacency matrix ([Wei]).

Proposition (Prop. 2.8)**.** *There exists a constant* $L > 0$ *such that for any* $r \geq 1$ *and any word* $w \in \mathcal{W}(\mathcal{A}(r))$ *of length* $n \leq p_1^2 \cdots p_r^2$*, we have* $\eta_r(w) > L \cdot e^{nh(\eta_r)}$*.*

This uniformity implies that the measure η on $\mathcal{A}$ which is the weak* limit of the measures η_r also satisfies a lower Gibbs bound. When combined with the ergodicity of η, which follows from its being the $\bar{d}$-limit of the ergodic measures η_r, this implies by a well-known argument that it is the only measure of maximal entropy on $\mathcal{A}$ (cf. 3.11).

The existence of the factor maps ϕ_r above, combined with results of Shields and Thouvenot on entropy zero × Bernoulli processes, yields the following clean description of the maximal entropy measure-preserving dynamical systems, i.e. the systems $\boldsymbol{\mathcal{A}}(r) := (\mathcal{A}(r), T, \eta_r)$ and $\boldsymbol{\mathcal{A}} := (\mathcal{A}, T, \eta)$.

Theorem (§3.4)**.** *We have natural isomorphisms* $\boldsymbol{\mathcal{A}}(r) = \mathbb{Z}_r \times \mathcal{B}_r$ *and* $\boldsymbol{\mathcal{A}} = \mathcal{K} \times \mathcal{B}_{sq}$*, where* $\mathcal{B}_r$ *resp.* $\mathcal{B}$ *is the Bernoulli system of entropy equal to that of* $\boldsymbol{\mathcal{A}}(r)$ *resp.* $\boldsymbol{\mathcal{A}}$*.*

This shows that the very natural system $\mathcal{A}$ exhibits an unusually tractable structure, in addition to providing a nontrivial concrete setting in which the properties of sofic systems, joinings and the $\bar{d}$-distance can be fruitfully applied.

I would like to thank my advisor Peter Sarnak for numerous helpful discussions.

2. Finitely many primes

2.1. The finite-level systems.

Let $\mathcal{A}$ be the squarefree flow as defined above, and for each prime p let

$$\mathcal{A}(p) = \left\{ y \in \Omega^{(2)} : |\pi_p(\mathrm{supp}(y))| < p^2 \right\}$$

and for $1 \leq k \leq p^2 - 1$ let

$$\mathcal{A}_k(p) = \left\{ y \in \Omega^{(2)} : |\pi_p(\mathrm{supp}(y))| = p^2 - k \right\}.$$

Here $\pi_p : \mathbb{Z} \to \mathbb{Z}/p^2\mathbb{Z}$ is the projection. Each $\mathcal{A}_k(p)$ is a Borel subset of $\mathcal{A}(p)$, and one can check that $\mathcal{A}_1(p)$ is open and dense in $\mathcal{A}(p)$. $\mathcal{A}_k(p)$ isn't quite invariant (consider $(1, 0, 0, 0, ...)$); we should rather consider the set

$$\left\{ y \in \Omega^{(2)} : |\pi_p(\mathrm{supp}(y))| = p^2 - k \text{ and } b(p^2) \cap \mathrm{supp}(y) \text{ is infinite whenever it's nonempty} \right\}$$

where $b(p^2)$ denotes the arithmetic progression $\{b + \ell p^2 : \ell \geq 0\}$ for $1 \leq b \leq p^2$. This set is Borel and invariant (but neither open nor closed) and properly contained in $\mathcal{A}_k(p)$, but Poincaré recurrence implies that their difference has absolute measure zero (i.e. has measure zero for any T-invariant measure on $\mathcal{A}(p)$), so we consider them to be the same unless otherwise stated.

Observe that for any p we have up to absolute measure zero

$$\mathcal{A}(p) = \coprod_{k=1}^{p^2-1} \mathcal{A}_k(p). \tag{2.1}$$

It's important to note that the $\mathcal{A}_k(p)$ are not compact, and so not suitable for studying the topological properties of $\mathcal{A}(p)$. However, they will be useful later when considering ergodic measures, where only the Borel structure is needed.

Proposition 2.1. *For any p we have*

$$h_{top}(\mathcal{A}(p)) = \left(1 - \frac{1}{p^2}\right) \log 2.$$

Proof. We use the following well-known formula for the topological entropy of a subshift $X \subset \Omega^{(2)}$ (it appears as Cor. 14.7 in [Gla] but is misstated there): letting u_n denote the number of distinct words $w = w_1 w_2 \cdots w_n$ of length n on the alphabet $\{0, 1\}$ that have form $w_i = x(i)$ for $i = 1, 2, ..., n$ for some $x \in X$, we have

$$h_{\text{top}}(X) = \lim_{n\to\infty} \frac{1}{n} \log u_n.$$

In our setting, a sequence $w \in \{0, 1\}^{mp^2}$ has form $w_i = y(i)$ for some $y \in \mathcal{A}(p)$ if and only if there exists some $a \in \left\{1, 2, ..., p^2\right\}$ (there may be more than one) such that $w_{a+\ell p^2} = 0$ for $\ell = 0, 1, 2, ..., m - 1$. The remaining entries of w may be chosen freely from $\{0, 1\}$. Hence, each choice a of an element of $\left\{1, 2, ..., p^2\right\}$ yields $2^{mp^2 - m} = 2^{m(p^2-1)}$ distinct admissible words of length $n = mp^2$. However, a word may be admissible for more than one choice of a, so we get the inequality

$$u_{mp^2} \leq p^2 2^{m(p^2-1)}$$

and therefore

$$\begin{aligned} h_{\text{top}}(\mathcal{A}(p)) &= \lim_{m\to\infty} \frac{1}{mp^2} \log u_{mp^2} \\ &\leq \lim_{m\to\infty} \frac{1}{mp^2} \left[\log\left(p^2 2^{m(p^2-1)}\right)\right] \\ &= \lim_{m\to\infty} \frac{m(p^2-1)}{mp^2} \log 2 \\ &= \left(1 - \frac{1}{p^2}\right) \log 2. \end{aligned}$$

But we also have the opposite inequality: since for a fixed choice a there are precisely $2^{m(p^2-1)}$ words of length mp^2 having only zeros along the length m arithmetic sequence modp^2 on a, we have

$$u_{mp^2} \geq 2^{m(p^2-1)}$$

from which we get $h_{\text{top}}(\mathcal{A}(p)) \geq (1 - 1/p^2) \log 2$, as desired.

□

Since the $\mathcal{A}_k(p)$ are disjoint and invariant, for each ergodic measure η on $\mathcal{A}(p)$ there exists a unique k such that $\eta(\mathcal{A}_k(p)) = 1$ while $\eta(\mathcal{A}_j(p)) = 0$ for $j \neq k$, and so $h_\eta(\mathcal{A}(p)) = h_\eta(\mathcal{C}_k(p))$ where $\mathcal{C}_k(p)$ is the closure of $\mathcal{A}_k(p)$.

Proposition 2.2. *For any prime p and $1 \leq k \leq p^2 - 1$, we have*

$$h_{top}(\mathcal{C}_k(p)) = \left(1 - \frac{k}{p^2}\right) \log 2.$$

Proof. For any invariant subset $X \subset \Omega^{(2)}$, the closure $\overline{X}$ is also invariant and has the same language as X, i.e. $\mathcal{W}(\overline{X}) = \mathcal{W}(X)$ ([Tho] pg. 3561). It then follows from the above proof that

$$h_{\text{top}}(\overline{X}) = \lim_{n\to\infty} \frac{1}{n} \log u_n$$

where u_n is the number of words of length n appearing in X. An easy extension of the argument in the above proposition shows that for $\mathcal{A}_k(p)$ we have the inequalities

$$2^{m(p^2-k)} \leq u_{mp^2} \leq \binom{p^2}{k} 2^{m(p^2-k)}$$

and since $\binom{p^2}{k}$ is independent of m, the desired result follows immediately. □

It follows from this and the proposition that if η is an ergodic measure on $\mathcal{A}(p)$, then η has maximal entropy if and only if its restriction to $\mathcal{A}_1(p)$ has maximal entropy (we're using the fact that $h_\eta(\mathcal{A}_k(p)) \leq h_{\text{top}}(\mathcal{C}_k(p))$ for any invariant measure η on $\mathcal{A}_k(p)$). In particular, η must be supported on $\mathcal{A}_1(p)$. Now suppose that η is any invariant (not necessarily ergodic) measure of maximal entropy on $\mathcal{A}(p)$. Then by Thm. 15.12 of [Gla] we have $h_\eta(\mathcal{A}(p)) = \int_Y h_{\eta_y}(\mathcal{A}(p))d\nu(y)$ where $\eta = \int_Y \eta_y d\nu(y)$ is the ergodic decomposition of η, so η_y is an *ergodic* measure of maximal entropy on $\mathcal{A}(p)$ for ν-a.e. $y \in Y$. The above then implies that η_y is supported on $\mathcal{A}_1(p)$ for ν-a.e. $y \in Y$, and it follows that η is supported on $\mathcal{A}_1(p)$ as well.

We now consider the situation for more than one prime. Let $p_1, ..., p_r$ be a finite collection of primes and set

$$\begin{aligned}\mathcal{A}(p_1, ..., p_r) &= \left\{y \in \Omega^2 : |\pi_{p_i}(\text{supp}(y))| < p_i^2, i = 1, 2, \dots, r\right\}\\ &= \cap_{i=1}^r \mathcal{A}(p_i).\end{aligned}$$

Also, for each r-tuple $\vec{k} = (k_i) \in \prod_{i=1}^r \{1, 2, ..., p_i^2 - 1\}$ define

$$\begin{aligned}\mathcal{A}_{\vec{k}}(p_1, \dots, p_r) &= \left\{y \in \Omega^2 : |\pi_{p_i}(\text{supp}(y))| = p_i^2 - k_i, i = 1, 2, \dots, r\right\}\\ &= \cap_{i=1}^r \mathcal{A}_{k_i}(p_i).\end{aligned}$$

We have the following obvious extension of Prop. 2.1.

Proposition 2.3. *For any $p_1, \dots, p_r$ and $\vec{k}$ as above, we have*

$$h_{top}(\mathcal{C}_{\vec{k}}(p_1, \dots, p_r)) = \log 2 \prod_{i=1}^r \left(1 - \frac{k_i}{p_i^2}\right)$$

where $\mathcal{C}_{\vec{k}}(p_1, \dots, p_r)$ is the closure of $\mathcal{A}_{\vec{k}}(p_1, \dots, p_r)$, and

$$h_{top}(\mathcal{A}(p_1, \dots, p_r)) = \max_{\vec{k}} h_{top}(\mathcal{C}_k(p_1, \dots, p_r)) = \log 2 \prod_{i=1}^r \left(1 - \frac{1}{p_i^2}\right).$$

Proof. This is an easy consequence of the above arguments, since by the Chinese remainder theorem and the inclusion-exclusion principle we have the inequalities

$$2^{m\prod_{i=1}^r (p_i^2 - k_i)} \leq u^{\vec{k}}_{mp_1^2 p_2^2 \cdots p_r^2} \leq \binom{p_1^2}{k_1}\binom{p_2^2}{k_2} \cdots \binom{p_r^2}{k_r} 2^{m\prod_{i=1}^r (p_i^2 - k_i)},$$

where $u^{\vec{k}}_{mp_1^2 p_2^2 \cdots p_r^2}$ is the number of distinct words of length $mp_1^2 p_2^2 \cdots p_r^2$ appearing in elements of $\mathcal{A}_{\vec{k}}(p_1, \dots, p_r)$. The conclusion follows at once. □

As per the remarks following Prop. 2.2, it follows from this that every invariant measure of maximal entropy on $\mathcal{A}(p_1, \ldots, p_r)$ is supported on $\mathcal{A}_{(1,1,\ldots,1)}(p_1, \ldots, p_r)$, which we abbreviate $\mathcal{A}_1(p_1, \ldots, p_r)$.

2.2. **Sofic systems.** Let X be a closed, invariant subset of the full 2-shift $(\Omega^{(2)}, T)$ and let $\mathcal{W}(X)$ be the set of words that appear in the elements of X. For each word $w \in \mathcal{W}(X)$ define

$$F(w) = \{v \in \mathcal{W}(X) : wv \in \mathcal{W}(X)\},$$

the so-called *follower set* of w. We consider F as a function $\mathcal{W}(X) \to 2^{\mathcal{W}(X)}$. The system (X, T) is then called *sofic* if the range of F is finite [Wei]. This is equivalent to the condition that X be a factor of a subshift of finite type. It is well-known that a topologically transitive subshift of finite type possesses a unique measure of maximal entropy; this was extended to sofic systems by Fischer and Weiss.

Theorem 2.4 ([Fis] Thm. 4)**.** *If (X, T) is sofic and topologically transitive, then it possesses a unique measure of maximal entropy.*

The construction of this measure is reviewed in the next subsection. A system possessing a unique measure of maximal entropy (which is then automatically ergodic) is called *intrinsically ergodic*. Our reason for introducing sofic systems is the following.

Proposition 2.5. *$\mathcal{A}(p)$ is sofic and topologically transitive. Hence, it is intrinsically ergodic by the above theorem.*

Proof. We begin by showing that $\mathcal{A}(p)$ is topologically transitive. Let $C_{A,B}, C_{A',B'} \subset \Omega^{(2)}$ be basic cylinders such that $U := C_{A,B} \cap \mathcal{A}(p)$ and $V := C_{A',B'} \cap \mathcal{A}(p)$ are both nonempty. We must show that there exists some $n \geq 0$ such that $U \cap T^{-n}V \neq \emptyset$. First, as A, B, A', B' are all finite, we may choose a large m such that $\max A < \min(B' + mp^2)$ and $\max B < \min(A' + mp^2)$. We then have for any $n \in \{0, 1, ..., p^2 - 1\}$ that

$$U \cap T^{-(n+mp^2)}V = C_{A \cup (A'+n+mp^2), B \cup (B'+n+mp^2)} \cap \mathcal{A}(p).$$

The intersection on the right will clearly be nonempty so long as the reduction $\bmod p^2$ of $A \cup (A'+n+mp^2)$, which equals the reduction $\bmod p^2$ of $A \cup (A'+n)$, is not all of $\mathbb{Z}/p^2\mathbb{Z}$; our task is thus reduced to finding an n for which this holds. Since U and V are nonempty it must be the case that $S = \pi_p(A)$ and $S' = \pi_p(A')$ are both proper subsets of $\mathbb{Z}/p^2\mathbb{Z}$. But then, if no such n existed, we would have $S \cup (S'+n) = \mathbb{Z}/p^2\mathbb{Z} \; \forall \, n \in \{0, 1, ..., p^2 - 1\}$. Choosing $x \notin S$, this would imply $x - n \in S' \; \forall \, n$, and therefore $|S'| = p^2$, a contradiction. Hence, $\mathcal{A}(p)$ is topologically transitive.

We now prove that $\mathcal{A}(p)$ is sofic. Let $\mathcal{W}$ be the set of all words on the alphabet $\{0, 1\}$ that appear in elements of $\mathcal{A}(p)$. For each $w = w_1 w_2 \cdots w_d \in \mathcal{W}$ (where $w_i \in \{0, 1\}$) with $d \geq p^2$, let $f = \lfloor d/p^2 \rfloor$ and define

$$S_p(w) = \{w_{ip^2+1} w_{ip^2+2} \cdots w_{(i+1)p^2} \in \mathcal{W} : i = 0, ..., f-1\} \cup \{w_{fp^2} \cdots w_d\}.$$

That is, $S_p(w)$ consists precisely of the distinct subblocks of w of length p^2 (as well as the final subblock of length $d - fp^2$). Clearly, $S_p(w)$ is finite, and moreover there are only finitely many possibilities for $S_p(w)$ as w varies over $\mathcal{W}$.

I claim that if $S_p(w) = S_p(w')$ for two words $w, w' \in \mathcal{W}$ both of length at least p^2 (but not necessarily of equal length), then $F(w) = F(w')$. It then follows from the above (and the fact that there are finitely many words in $\mathcal{W}$ of length less than p^2) that $\mathcal{A}(p)$ is sofic. To prove the claim, suppose $\mathcal{S}_p(w) = \mathcal{S}_p(w')$ and let $v = v_1 \cdots v_r \in F(w)$. We wish to show that $w'v \in \mathcal{W}$, and thus that $v \in F(w')$; as this is completely symmetric in w and w', this implies $F(w) = F(w')$. Now, suppose that $w'\upsilon \notin \mathcal{W}$. Then there exist $\ell_0, \ell_1, \ell_2, ..., \ell_{p^2-1} \geq 0$ such that

$$(w'\upsilon)_{a+\ell_a p^2} = 1 \ \forall \ a \in \left\{0, 1, 2, ..., p^2 - 1\right\}.$$

Since w' and υ both belong to $\mathcal{W}$, there is a partition $\left\{0, 1, 2, ..., p^2 - 1\right\} = E \coprod F$ with E and F both nonempty, such that for all $a \in E$ we have $a + \ell_a p^2 \leq d'$ (d' is the length of w') and $w'_{a+\ell_a p^2} = 1$, while for all $b \in F$ we have $d' + 1 \leq b + \ell_b p^2 \leq d' + r$ and $\upsilon_{b+\ell_b p^2 - d'} = 1$.

Since $S_p(w') = S_p(w)$, it follows that for each $a \in E$ there exists $t_a \geq 0$ such that $w_{a+t_a p^2} = 1$. Thus, if we consider a sequence $y \in \mathcal{A}(p)$ such that $(w\upsilon)_j = y(j)$ for $j = 1, 2, ..., d + r$ (such a y exists because $w\upsilon \in \mathcal{W}$ and $\mathcal{A}(p)$ is shift invariant), then we have $y(a + t_a p^2) = 1 \ \forall \ a \in E$. For $b \in F$, since $d' + 1 \leq b + \ell_b p^2 \leq d' + r$ we have $d + 1 \leq d - d' + b + \ell_b p^2 \leq d + r$ and

$$\begin{aligned} y(d - d' + b + \ell_b p^2) &= (w\upsilon)_{d-d'+b+\ell_b p^2} \\ &= \upsilon_{b+\ell_b p^2 - d'} \ \text{ since } d - d' + b + \ell_b p^2 > d \\ &= 1 \end{aligned}$$

by the above. However, the fact that $S_p(w) = S_p(w')$ implies that $d \equiv d' \bmod p^2$, so $d - d' + b + \ell_b p^2 \equiv b \bmod p^2$. Therefore, $\mathrm{supp}(y)$ contains the sets $\{a + t_a p^2 : a \in E\}$ and $\{d - d' + b + \ell_b p^2 : b \in F\}$, and moreover the reduction $\bmod p^2$ of the union of these sets is $E \cup F = \left\{0, 1, 2, \ldots, p^2 - 1\right\}$. But this contradicts the fact that $y \in \mathcal{A}(p)$. ☐

In fact, the above proof shows that two words w, u appearing in $\mathcal{A}(p)$ have the same follower set if and only if $\pi_p(\mathrm{supp}(w)) = \pi_p(\mathrm{supp}(u))$ and $|w| \equiv |u| \bmod p^2$ (where $|w|$ is the length of w), or if w and u both consist entirely of zeros.

Proposition 2.6. *$\mathcal{A}(p_1, \ldots, p_r)$ is sofic and topologically transitive for any finite collection of primes $p_1, \ldots, p_r$, and therefore is intrinsically ergodic.*

Proof. $\mathcal{A}(p_1, \ldots, p_r)$ is sofic because it is the intersection of the sofic shifts $\mathcal{A}(p_i)$ for $i = 1, \ldots, r$, and the intersection of two sofic shifts is again sofic ([LM] Prop. 3.4.10). To prove that it is topologically transitive, let A, B, A', B' be finite subsets of $\mathbb{N}$ such that $U := C_{A,B} \cap \mathcal{A}(p_1, \ldots, p_r)$ and $V := C_{A',B'} \cap \mathcal{A}(p_1, \ldots, p_r)$ are nonempty. Then for each i, the reductions $S_i := \pi_{p_i}(A)$ and $S'_i := \pi_{p_i}(A')$ are proper subsets of $\mathbb{Z}/p_i^2\mathbb{Z}$. Hence, by the proof of Prop. 2.5, we may find for each i an $n_i \in \{0, 1, \ldots, p_i^2 - 1\}$ such that $S_i \cup (S'_i + n_i)$ is again proper. By the Chinese remainder theorem, there exists $n \in \mathbb{N}$ such that $n \equiv n_i \bmod p_i^2$ for $i = 1, \ldots, r$. Then, choosing a large m such that $\max A < \min(B' + mp_1^2 \cdots p_r^2)$ and $\max B < \min(A' + mp_1^2 \cdots p_r^2)$, the set

$$U \cap T^{-(n+mp_1^2 \cdots p_r^2)} V = C_{A \cup (A' + n + mp_1^2 \cdots p_r^2), B \cup (B' + n + mp_1^2 \cdots p_r^2)} \cap \mathcal{A}(p_1, \ldots, p_r)$$

is nonempty, as desired. ☐

2.3. **Properties of the measures of maximal entropy.** In this section we study the unique measure of maximal entropy η_r on $\mathcal{A}(r) := \mathcal{A}(p_1, \ldots, p_r)$. We will develop explicit formulas describing the behavior of these measures in order to derive properties of the full squarefree flow via a limit process.

Corresponding to the irreducible (= topologically transitive) sofic system $\mathcal{A}(r)$ are two finite directed labelled graphs G_r, G'_r both of whose vertex sets equal the set of equivalence classes of finite words appearing in $\mathcal{A}(r)$, where two words are considered equivalent if and only if their follower sets are equal. Observe that by the proof of Prop.



FIGURE 1. Graph presenting the sofic shift $\mathcal{A}(p)$ for $p = 2$.

2.5, two words $u, v \in \mathcal{W}(\mathcal{A}(r))$ are equivalent if and only if $|u| \equiv |v| \bmod \left(p_1^2 \cdots p_r^2\right)$ and $\pi_{p_i}(\mathrm{supp}(u)) = \pi_{p_i}(\mathrm{supp}(v))$ for $i = 1, 2, \ldots, r$, or if u and v both consist entirely of zeros. Therefore, the equivalence classes of words in $\mathcal{A}(r)$ (and therefore the vertices of both of the graphs G_r, G_r') are in bijection with pairs $(d, (S_i)_{i=1}^r)$ where $d \in \mathbb{Z}/p_1^2 \cdots p_r^2\mathbb{Z}$ and each S_i is a proper subset of $\mathbb{Z}/p_i^2\mathbb{Z}$, and the S_i are either all empty (corresponding to words consisting of only zeros) or all nonempty.

Letting $[u], [v]$ be vertices of G_r, there is a directed edge from $[u]$ to $[v]$ labelled 0 if and only if $[u0] = [v]$, and there is an edge labelled 1 if and only if $[u1] = [v]$. Similarly, there is a directed edge in G_r' from $[u]$ to $[v]$ labelled $i \in \{0, 1\}$ if and only if $[iu] = [v]$. To illustrate, figure 1 shows a piece of G_r in the case where we consider only the single prime $p = 2$. We don't draw the whole graph as it has $2^2(2^4 - 1) = 60$ vertices.

In this graph, we've made use of the fact that, for instance, $[011001] = [011000]$. Indeed, the words 011001 and 011000 both have length congruent to 2 mod 4 and the reduction mod4 of both their supports is the same set, namely $\{2, 3\}$ mod 4. Observe that a sequence of zeros and ones belongs to $\mathcal{A}(r)$ if and only if it can be obtained by reading in order the labels of the edges in an infinite directed walk on G_r. This is the same as saying that $\mathcal{A}(r)$ is recognizable by the finite automaton G_r (as is true of any sofic system; see [LM]).

Define the *period* of $\mathcal{A}(r)$ to be the greatest common divisor of the lengths of the cycles in the graph $G(r)$ (this definition makes sense for any sofic system). We then have the following easy observation, which will be needed later.

Proposition 2.7. *$\boldsymbol{\mathcal{A}}(r)$ has period $p_1^2 \cdots p_r^2$.*

Proof. We first prove that $\boldsymbol{\mathcal{A}}(p)$ has period p^2. The graph G_p defining this system was described in section 2.3. This graph need not be irreducible, but $\mathcal{A}(p)$ is presented by one of its maximal irreducible components, so we make no distinction between them.

One can see from the description of G_p that if a path $c = i_1 i_2 \cdots i_n$ in G_p begins and ends at the same vertex $[w]$ (i.e. c is a cycle), then we must have $|w| \equiv |w| + n (\bmod p^2)$, and therefore $p^2 | n$. Conversely, if we choose a vertex $[w]$ and create a directed path in G_p by beginning at $[w]$ and travelling only along edges labelled 0, then this path will return to $[w]$ after exactly p^2 steps. Therefore $\boldsymbol{\mathcal{A}}(p)$ has period p^2.

We can deduce directly from this that $\boldsymbol{\mathcal{A}}(r)$ has period $p_1^2 \cdots p_r^2$. By [LM] Thm. 3.4.10, the graph representing a finite intersection of sofic shifts is simply the label product of the graphs representing each of the shifts, and it's an easy exercise to show that the period of the label product of two irreducible graphs is the least common multiple of their periods, from which the claim follows. $\square$

Let $T(r)$ resp. $T'(r)$ be the adjacency matrix of the finite directed graph G_r resp. G'_r, so $T(r)_{[u][v]}$ is the number of directed edges in G_r which begin at the vertex $[u]$ and end at the vertex $[v]$ (and similarly for $T'(r)$). By the proof of Thm. 1 of [Wei] we have the following facts: letting $\lambda_r = e^{h_{\text{top}}(\mathcal{A}(r))}$, λ_r is an eigenvalue of both $T(r)$ and $T'(r)$. We have $T(r)T'(r) = T'(r)T(r)$ and the matrix $F(r) := T(r)T'(r)$ is irreducible, with Perron-Frobenius eigenvalue λ_r^2. There exist strictly positive right and left eigenvectors of $F(r)$ both with eigenvalue λ_r^2, which are unique up to multiplication by positive real numbers. Moreover, they're eigenvectors of both $T(r)$ and $T'(r)$, with eigenvalue λ_r in both cases.

A measure on $\mathcal{A}(r)$ is completely determined by its behavior on the cylinder sets

$$\mathcal{A}(r)_w = \{x \in \mathcal{A}(r) : x(i) = w_i \text{ for } i = 1, \dots, n\}$$

where w is a length n word appearing in $\mathcal{A}(r)$. The measure of maximal entropy on $\mathcal{A}(r)$ is then given by

$$\eta_r \left(\mathcal{A}(r)_w\right) = \lambda_r^{-n} m(r)_{[w]} \tag{2.2}$$

where $m(r)$ is the strictly positive right Perron-Frobenius eigenvector for $F(r)$, normalized so that $m(r)_{[0]} + m(r)_{[1]} = \lambda_r$ (cf. [Wei] Eqn. (14)). Our aim is to make this expression concrete enough that we can consider the limiting behavior of these measures when w is a fixed word in the full squarefree flow $\mathcal{A}$ and $r \to \infty$. Such considerations allow us to prove in section 3.3 that $\mathcal{A}$ is intrinsically ergodic.

We begin by considering a single prime p; the limit as we include more primes is of course handled using the Chinese remainder theorem. Let w be an arbitrary word appearing in $\mathcal{A}$ and define the cylinder set $\mathcal{A}(p)_w$ as above. The idea is that since η_p is supported on $\mathcal{A}_1(p)$ by the remarks following Prop. 2.2, we should replace w by extensions of it that omit only one residue class (we assume $p^2 \gg |w|$ so this can't already be the case). Indeed, let $a_1, \dots, a_k$ be the classes modp^2 omitted by supp(w). Then Prop. 2.2 implies that up to η_p-measure zero we have

$$\begin{aligned}
\mathcal{A}(p)_w &= \{x \in \mathcal{A}_1(p) : w \subset x\} \\
&= \coprod_{i=1}^{k} \left\{x \in \mathcal{A}_1(p) : w \subset x \text{ and } \pi_p(\text{supp}(x)) = \left(\mathbb{Z}/p^2\mathbb{Z}\right) \backslash \{a_i\}\right\} \\
&= \coprod_{i=1}^{k} \coprod_{m=1}^{\infty} \coprod_{\substack{w \subset w' \\ |w'| = p^2 \\ \pi_p(\text{supp}(w')) = (\mathbb{Z}/p^2\mathbb{Z}) \backslash \{a_i\}}} \{x \in \mathcal{A}(p) : w' \subset x \text{ and } m(x) = m\}.
\end{aligned}$$

Here, $m(x) = \min\{m : \pi_p(\text{supp}(x_1 \cdots x_{mp^2})) = \pi_p(\text{supp}(x))\}$, and the notation $w \subset x$ means $x(1) \cdots x(|w|) = w$ (and similarly for $w \subset w'$). The definition of $m(x)$ as a minimum is needed to produce a disjoint union in the above decomposition. Define

$$C^p_m(w,i) = \{w' \in \mathcal{W}(\mathcal{A}(p)) : w \subset w', |w'| = mp^2, \pi_p(\mathrm{supp}(w')) = \mathbb{Z}/p^2\mathbb{Z}\backslash\{a_i\} \text{ and } |\pi_p(\mathrm{supp}(w'_1 \cdots w'_{np^2}))| < p^2 - 1 \text{ for } n < m\}$$

Then the above implies that up to η_p-measure zero

$$\mathcal{A}(p)_w = \coprod_{i=1}^{k} \coprod_{m=1}^{\infty} \coprod_{w' \in C^p_m(w,i)} \mathcal{A}(p)_{w'}.$$

Therefore,

$$\eta_p(\mathcal{A}(p)_w) = \sum_{i=1}^{k} \sum_{m=1}^{\infty} \sum_{w' \in C^p_m(w,i)} \eta_p(\mathcal{A}(p)_{w'})$$

and since any word in $C^p_m(w,i)$ is equivalent to the word $\nu_i = 11\cdots101\cdots1$ which has length p^2 and where the 0 occurs at a_i, 2.2 gives

$$\begin{aligned}\eta_p(\mathcal{A}(p)_w) &= \sum_{i=1}^{k} \sum_{m=1}^{\infty} \sum_{w' \in C^p_m(w,i)} \lambda_p^{-mp^2} m(p)_{[\nu_i]} \\ &= \sum_{i=1}^{k} m(p)_{[\nu_i]} \sum_{m=1}^{\infty} \lambda_p^{-mp^2} \left|C^p_m(w,i)\right|.\end{aligned}$$

To determine $|C^p_m(w,i)|$, observe that if $w' \in C^p_m(w,i)$, the condition $|\pi_p(\mathrm{supp}(w'_1 \cdots w'_{np^2}))| < p^2 - 1$ for $n < m$ in the definition of $C^p_m(w,i)$ implies that there is a $j \neq i$ such that $w'_{a_j+\ell p^2} = 0$ for $1 \leq \ell < m-1$ while $w'_{a_j+(m-1)p^2} = 1$. Thus the $m-2$ positions $a_j + \ell p^2, 1 \leq \ell < m-1$ of w' are fixed to be 0; since the position $a_j + (m-1)p^2$ is 1, there are at most $mp^2 - n - (m-1) - m$ positions in w' to be freely filled (taking into account $w \subset w'$) once a_j is chosen from the $k-1$ possibilities, where the second m comes from w' having to omit $a_i(p^2)$. However, since the reduction of the support of w' is $(\mathbb{Z}/p^2\mathbb{Z})\backslash\{a_i\}$, w' must also have a 1 somewhere in each arithmetic sequence $a_s(p^2)$ for $s \neq i, j$. As there are m choices of a position for such a 1 for each $s \neq i, j$, and there are $k-2$ such s, we see that

$$|C^p_m(w,i)| = (k-1)m^{k-2}2^{m(p^2-1)-n-(m-1)-(k-2)}$$

(the factor of $k-1$ in front comes from choosing a_j) and therefore (assuming $k > 1$)

$$\eta_p(\mathcal{A}(p)_w) = (k-1) \sum_{i=1}^{k} m(p)_{[\nu_i]} \sum_{m=1}^{\infty} \lambda_p^{-mp^2} m^{k-2} 2^{m(p^2-1)-n-(m-1)-(k-2)}.$$

Now,

$$\begin{aligned}\lambda_p^{-mp^2} &= \left(2^{(1-1/p^2)}\right)^{-mp^2} \\ &= 2^{-m(p^2-1)}\end{aligned}$$

so we get

$$\eta_p(A(p)_w) = (k-1)2^{-n-k+3} \sum_{i=1}^{k} m(p)_{[\nu_i]} \sum_{m=1}^{\infty} m^{k-2}2^{-m}. \tag{2.3}$$

The sum $\sum_{m=1}^{\infty} m^{k-2}2^{-m}$ clearly converges to a value $c(k)$ independent of i; since k depends on p and w, so does $c(k)$.

To analyze the terms $m(p)_{[\nu_i]}$, we use the strictly positive left Perron-Frobenius eigenvector $m'(p)$ for $F(p)$, normalized so that $m(p) \cdot m'(p) = 1$. It's related to the right eigenvector by the formula ([KiS] pg. 235)

$$m(p)_{[\nu]} m'(p)_{[\nu]} = \lim_{n\to\infty} \frac{(F(p)^{np^2})_{[\nu][\nu]}}{\lambda_p^{2np^2}} \tag{2.4}$$

for $[\nu]$ belonging to the maximal irreducible component of G_p. The factor p^2 appears because it's the period of $\mathcal{A}(p)$ by Prop. 2.7 above, while the 2 in the exponent of λ_p comes from the fact that λ_p^2, not λ_p, is an eigenvalue of $F(p)$.

In our setting, the maximal irreducible component of G_p is the induced subgraph whose vertices consist of the equivalence classes of words in $\mathcal{A}(p)$ whose support omits only one residue class modp^2. If $[\nu]$ is such an equivalence class, then we see from the definition of $F(p)$ and the graphs G_p, G'_p that $(F(p)^{np^2})_{[\nu][\nu]}$ equals the number of sequences of pairs $(i_1, j_1), \ldots, (i_{np^2}, j_{np^2})$ such that $[j_{np^2} \cdots j_1 \nu i_1 \cdots i_{np^2}] = [\nu]$. Since ν omits only one residue class $\bmod p^2$, we're free to choose such a sequence of pairs so long as we get zeros along the length $2np^2$ arithmetic sequence on the residue class omitted by ν, which determines exactly $2n$ positions in the word $j_{np^2} \cdots j_1 \nu i_1 \cdots i_{np^2}$. Hence, the total number of such sequences of pairs is $2^{2n(p^2-1)}$. Since we also have $\lambda_p^{2np^2} = 2^{2n(p^2-1)}$, (2.4) gives $m(p)_{[\nu]} m'(p)_{[\nu]} = 1$. (2.3) therefore becomes

$$\eta_p(\mathcal{A}(p)_w) = (k-1) 2^{-n-k+3} c(k) \sum_{i=1}^{k} m'(p)^{-1}_{[\nu_i]}. \tag{2.5}$$

To illustrate the idea of our analysis, we calculate $m'(p)_{[11\cdots 10]}$ (note that this need not appear in the above sum). The eigenvector equation $m'(p)T(p) = \lambda_p m'(p)$ gives for any state $[\nu]$

$$m'(p)_{[\nu]} = \lambda_p^{-1} \sum_{[\nu']} n(\nu', \nu) m'(p)_{[\nu']} \tag{2.6}$$

where the sum is over all equivalence classes of words in $\mathcal{A}(p)$ and $n(\nu', \nu) = |\{i \in \{0,1\} : [\nu' i] = [\nu]\}|$. This immediately gives

$$m'(p)_{[11\cdots 10]} = \lambda_p^{-1} m'(p)_{[v_1]}$$

where v_1 is the word of length $p^2 - 1$ consisting only of ones. For $0 \le i \le p^2$, let v_i be the word consisting of $p^2 - i$ ones, and let u_i be the word consisting of $p^2 - i$ zeros; then we have $[v_1] = [v_2 1]$ and also $[v_1 u_1 0] = [v_1] = [v_1 u_1 1]$. Therefore

$$\begin{aligned} m'(p)_{[11\cdots 10]} &= \lambda_p^{-1} m'(p)_{[v_1]} \\ &= \lambda_p^{-2} (m'(p)_{[v_2]} + 2m'(p)_{[v_1 u_1]}). \end{aligned} \tag{2.7}$$

Again by (2.6), we have

$$\begin{aligned} m'(p)_{[v_1 u_1]} &= \lambda_p^{-1} m'(p)_{[v_1 u_2]} \\ &= \cdots \\ &= \lambda_p^{-(p^2-1)} m'(p)_{[v_1]} \end{aligned}$$

so that, by (2.7),

$$\lambda_p^{-1} m'(p)_{[v_1]} = \lambda_p^{-2} \left(m'(p)_{[v_2]} + 2\lambda_p^{-(p^2-1)} m'(p)_{[v_1]} \right)$$

and solving for $m'(p)_{[v_1]}$ yields

$$\begin{aligned} m'(p)_{[11\cdots 10]} &= \lambda_p^{-1} m'(p)_{[v_1]} \\ &= \lambda_p^{-2}(1-2\lambda_p^{-p^2})^{-1} m'(p)_{[v_2]}. \end{aligned}$$

Similarly, since

$$m'(p)_{[v_2]} = \lambda_p^{-1}\left(m'(p)_{[v_3]} + 2m'(p)_{[v_2 u_1]}\right)$$

analyzing as above yields

$$m'(p)_{[v_2]} = \lambda_p^{-1}(1-2\lambda_p^{-p^2})^{-1} m'(p)_{[v_3]}$$

and so

$$\begin{aligned} m'(p)_{[11\cdots 10]} &= \lambda_p^{-2}(1-2\lambda_p^{-p^2})^{-1} m'(p)_{[v_2]} \\ &= \lambda_p^{-3}(1-2\lambda_p^{-p^2})^{-2} m'(p)_{[v_3]}. \end{aligned}$$

Continuing this procedure, we find the formula

$$m'(p)_{[11\cdots 10]} = \lambda_p^{-(p^2-1)}(1-2\lambda_p^{-p^2})^{-(p^2-2)} m'(p)_{[1]}.$$

Analyzing the terms $m'(p)^{-1}_{[\nu_i]}$ in (2.5) in the same way gives exactly the same result, and so we get (assuming w doesn't omit 1 mod p^2)

$$\begin{aligned} \eta_p(A(p)_w) &= (k-1)2^{-n-k+3}c(k)\sum_{i=1}^{k} m'(p)^{-1}_{[\nu_i]} \\ &= 2^{-n-k+3}(k-1)kc(k)\lambda_p^{p^2-1}\left(1-2\lambda_p^{-p^2}\right)^{p^2-2} m'(p)^{-1}_{[1]}. \end{aligned} \tag{2.8}$$

We now consider the case when $r > 1$. For $i = 1,\ldots,r$, let C_i be the set of residue classes modp_i^2 omitted by w. Then as above, Prop. 2.3 implies that up to η_r-measure zero we have

$$\begin{aligned} A(r)_w &= \{x \in \mathcal{A}_1(r) : w \subset x\} \\ &= \coprod_{(a^{(1)},a^{(2)},\ldots,a^{(r)})\in\prod_{i=1}^r C_i} \left\{x \in \mathcal{A}_1(r) : w \subset x \text{ and } \pi_{p_i}(\mathrm{supp}(x)) = \left(\mathbb{Z}/p_i^2\mathbb{Z}\right)\setminus\{a^{(i)}\}\right\} \\ &= \coprod_{\substack{(a^{(1)},a^{(2)},\ldots,a^{(r)}) \\ \in\prod_{i=1}^r C_i}} \coprod_{m=1}^{\infty} \coprod_{\substack{w\subset w' \\ |w'|=mp_1^2\cdots p_r^2 \\ \pi_{p_i}(\mathrm{supp}(w'))=\left(\mathbb{Z}/p_i^2\mathbb{Z}\right)\setminus\{a^{(i)}\}}} \{x \in \mathcal{A}(r) : w' \subset x \text{ and } m(x) = m\} \end{aligned}$$

where, as in the case of a single prime, $m(x) = \min\{m : \pi_{p_i}(\mathrm{supp}(x_1\cdots x_{mp_1^2\cdots p_r^2})) = \pi_{p_i}(\mathrm{supp}(x))$ for $i = 1,\ldots,r\}$. For each tuple $a = (a^{(1)},a^{(2)},\ldots,a^{(r)}) \in \prod_{i=1}^r C_i$ and $m \geq 1$ define

$$\begin{aligned} C_m^r(w,a) = \Big\{ w' \in \mathcal{W}(\mathcal{A}(r)) : w \subset w', |w'| = mp_1^2\cdots p_r^2, \pi_{p_i}(\mathrm{supp}(w')) = \left(\mathbb{Z}/p_i^2\mathbb{Z}\right)\setminus\{a^{(i)}\} \\ \text{for } i = 1,\ldots,r \text{ and, for } n < m, \exists i \text{ such that} \\ |\pi_{p_i}(\mathrm{supp}(w'_1\cdots w'_{np_1^2\cdots p_r^2}))| < p_i^2 - 1\Big\}. \end{aligned}$$

Then by the above, we have up to η_r-measure zero that

$$\eta_r(A(r)_w) = \sum_{a\in\prod_{i=1}^r C_i}\sum_{m=1}^{\infty}\sum_{w'\in C_m^r(w,a)} \eta_r(A(r)_{w'})$$

and since any word in $C_m^r(w,a)$ is equivalent to the word $\nu_r(a)$ of length $p_1^2\cdots p_r^2$ which has zeros along each arithmetic sequence $a^{(i)}(p_i^2)$ and ones everywhere else, (2.2) gives

$$\eta_r(A(r)_w) = \sum_{a\in\prod_{i=1}^r C_i} m(r)_{[\nu_r(a)]} \sum_{m=1}^{\infty} \lambda_r^{-mp_1^2\cdots p_r^2} |C_m^r(w,a)|.$$

We now give a lower bound for $|C_m^r(w,a)|$ that will be useful later. Observe that the last condition in the definition of $C_m^r(w,a)$ is satisfied for any $n < m$ as soon as it's satisfied for $n = m-1$. Choosing a $j \le r$ and $b^{(j)} \neq a^{(j)}$ a residue class $\mathrm{mod} p_j^2$ to be omitted by $w_1'\cdots w_{(m-1)p_1^2\cdots p_r^2}'$ specifies $(m-1)p_1^2\cdots \widehat{p_j^2}\cdots p_r^2$ positions in w' to be zero in addition to the $m\prod_{i=1}^r(p_i^2-1)+n$ positions specified by the other conditions in the definition of $C_m^r(w,a)$. So we find

$$|C_m^r(w,a)| \ge \max_{j=1,\dots,r} 2^{mp_1^2\cdots p_r^2 - m\prod_{i=1}^r(p_i^2-1)-(m-1)p_1^2\cdots\widehat{p_j^2}\cdots p_r^2-n}.$$

Since $\lambda_r^{-mp_1^2\cdots p_r^2} = 2^{-m\prod_{i=1}^r(p_i^2-1)}$, we get

$$\begin{aligned}\eta_r(A(r)_w) &\ge 2^{-n} \sum_{a\in\prod_{j=1}^r C_j} m(r)_{[\nu_r(a)]}\\ &\qquad\times \sum_{m=1}^{\infty} \lambda_r^{-mp_1^2\cdots p_r^2} \max_{j=1,\dots,r} 2^{mp_1^2\cdots p_r^2 - m\prod_{i=1}^r(p_i^2-1)-(m-1)p_1^2\cdots\widehat{p_j^2}\cdots p_r^2}\\ &\ge 2^{-n} \max_{j=1,\dots,r} 2^{p_1^2\cdots\widehat{p_j^2}\cdots p_r^2} s_r(j) \sum_{a\in\prod_{j=1}^r C_j} m(r)_{[\nu_r(a)]} \qquad (2.9)\end{aligned}$$

where $s_r(j) = \sum_{m=1}^{\infty} 2^{m\left[p_1^2\cdots p_r^2 - 2\prod_{i=1}^r(p_i^2-1)-p_1^2\cdots\widehat{p_j^2}\cdots p_r^2\right]}$.

We now turn to the terms $m(r)_{[\nu_r(a)]}$. We again use the strictly positive left Perron-Frobenius eigenvector $m(r)$ of $F(r)$; just as with (2.4) we have

$$\begin{aligned}m(r)_{[\nu]}m'(r)_{[\nu]} &= \lim_{n\to\infty} \frac{(F(r)^{np_1^2\cdots p_r^2})_{[\nu][\nu]}}{\lambda_r^{2np_1^2\cdots p_r^2}}\\ &= 1. \qquad (2.10)\end{aligned}$$

To illustrate the analysis of $m'(r)_{[\nu_r(a)]}$, we consider the case when $a = (0,0,\dots,0)$, so $\nu_r(a) =: \nu$ is the word of length $p_1^2\cdots p_r^2$ having zeros at the multiples of p_i^2 for each i and ones everywhere else. We proceed as we did with $m'(p)_{[\nu_i]}$ above. For each $s \ge 0$ let ν_s be the first subword of ν of length $p_1^2\cdots p_r^2 - s$. We have

$$m'(r)_{[\nu]} = \lambda_r^{-1}(m'(r)_{[\nu_1]} + 2m'(r)_{[\nu u_1]})$$

where u_1 is the word consisting of $p_1^2\cdots p_r^2-1$ zeros, and more generally u_s is the word consisting of $p_1^2\cdots p_r^2 - s$ zeros. Now

$$\begin{aligned}m'(r)_{[\nu u_1]} &= \lambda_r^{-1}m'(r)_{[\nu u_2]}\\ &= \lambda_r^{-2}m'(r)_{[\nu u_3]}\\ &= \cdots\\ &= \lambda_r^{-p_1^2\cdots p_r^2-1}m'(r)_{[\nu]}\end{aligned}$$

and so

$$m'(r)_{[\nu]} = \lambda_r^{-1}(1-2\lambda_r^{-p_1^2\cdots p_r^2})^{-1}m'(r)_{[\nu_1]}.$$

Repeating this for $m'(r)_{[\nu_1]}$, we find as with $m'(p)_{[\nu_i]}$

$$m'(r)_{[\nu]} = \lambda_r^{-p_1^2}(1-2\lambda_r^{-p_1^2\cdots p_r^2})^{-p_1^2} m'(r)_{[\nu_{p_1^2}]}.$$

We've stopped after p_1^2 steps because the last letter of $\nu_{p_1^2}$ is zero (which is not the case of any ν_s for $s < p_1^2$), so that

$$m'(r)_{[\nu_{p_1^2}]} = \lambda_r^{-1} m'(r)_{[\nu_{p_1^2+1}]}$$

and therefore

$$m'(r)_{[\nu]} = \lambda_r^{-p_1^2-1}(1-2\lambda_r^{-p_1^2\cdots p_r^2})^{-p_1^2} m'(r)_{[\nu_{p_1^2+1}]}.$$

We continue in this fashion, gaining a factor of λ_r^{-1} at each step and a factor of $(1-2\lambda_r^{-p_1^2\cdots p_r^2})^{-p_1^2}$ at step s if s not a multiple of p_i^2 for any i. The number of such s is precisely $\prod_{i=1}^r (p_i^2-1)$, so we find

$$m'(r)_{[\nu]} = \lambda_r^{-p_1^2\cdots p_r^2+1}(1-2\lambda_r^{-p_1^2\cdots p_r^2})^{-\prod_{i=1}^r(p_i^2-1)+1} m'(r)_{[1]}.$$

The same formula holds for $m'(r)_{[\nu_r(a)]}$ for any a, so (2.9) becomes

$$\eta_r(A(r)_w) \geq k_1\cdots k_r 2^{-n}\lambda_r^{p_1^2\cdots p_r^2-1}(1-2\lambda_r^{-p_1^2\cdots p_r^2})^{\prod_{i=1}^r(p_i^2-1)-1} m'(r)_{[1]}^{-1}\sum_{i=1}^r (k_i-1)2^{-k_i+2}s_r(k_i). \tag{2.11}$$

We now use this formula to provide a uniform lower bound for the entries $m(r)_{[w]}$ of the Perron-Frobenius eigenvector. Such a bound is crucial for proving in the next section that $\mathcal{A}$ carries a unique measure of maximal entropy. This reflects the importance of the rate of convergence (suitably interpreted) of the finite-level systems $\mathcal{A}(r)$ in establishing properties of the full squarefree flow $\mathcal{A}$. Indeed, any irreducible closed subshift whatsoever can be written canonically as a decreasing intersection of decreasing irreducible subshifts of finite type. Each of these is necessarily intrinsically ergodic, yet the intersection can carry any number of measures of maximal entropy (cf. [Hay]) since the systems need not converge in a uniform way. For another perspective on the way in which the rate of convergence of a decreasing sequence of intrinsically ergodic systems influences the properties of the limit, see for instance [Gur].

Proposition 2.8. *There exists a constant $L > 0$ such that for any $r \geq 1$ and any word $w \in \mathcal{W}(\mathcal{A}(r))$ of length at most $p_1^2\cdots p_r^2$, we have $m(r)_{[w]} > L$. In particular, if $w \in \mathcal{W}(\mathcal{A})$ (so that $w \in \mathcal{W}(\mathcal{A}(r))$ for any $r \geq 1$) we have $\liminf_{r\to\infty} m(r)_{[w]} \geq L$.*

Proof. Eq. (2.11) above states that if w is a word in $\mathcal{A}(r)$ of length $n \leq p_1^2\cdots p_r^2$ which omits k_i residue classes modp_i^2 for $i = 1,\ldots,r$ then

$$\eta_r(A(r)_w) \geq k_1\cdots k_r 2^{-n}\lambda_r^{p_1^2\cdots p_r^2-1}(1-2\lambda_r^{-p_1^2\cdots p_r^2})^{\prod_{i=1}^r(p_i^2-1)-1} m'(r)_{[1]}^{-1}\sum_{i=1}^r (k_i-1)2^{-k_i+2}s_r(k_i).$$

Here $s_r(k_i) = \sum_{m=1}^\infty (mp_1^2\cdots \widehat{p_i^2}\cdots p_r^2-1)^{k_i-2}2^{-mp_1^2\cdots\widehat{p_i^2}\cdots p_r^2}$. It can easily be verified that $2^{-k_i+2}s_r(k_i)$ is increasing with k_i so long as $k_i \geq 2$. Therefore, since $k_i \geq 1$ for each i and $\eta_r(w) = \lambda_r^{-n}m(r)_{[w]}$ we have the lower bound

$$m(r)_{[w]} \geq \lambda_r^{n+p_1^2\cdots p_r^2-1}2^{-n}(1-2\lambda_r^{-p_1^2\cdots p_r^2})^{\prod_{i=1}^r(p_i^2-1)-1} m'(r)_{[1]}^{-1}\sum_{i=1}^r s_r(k_i=2).$$

By eq. (2.10) we have $m'(r)_{[1]}^{-1} = m(r)_{[1]}$. Since the word $w = 1$ of course appears in $\mathcal{A}$, the weak* convergence of the η_r and eq. (2.2) show that the sequence $m(r)_{[1]}$ converges to $\lambda \cdot \eta(\mathcal{A}_1)$. In particular the sequence is bounded below by a positive number $m > 0$.

We have on letting $P_r = \prod_{i=1}^r (p_i^2 - 1)$

$$s_r(j) = \sum_{m=1}^{\infty} 2^{m\left[p_1^2 \cdots p_r^2 - 2P_r - p_1^2 \cdots \widehat{p_j^2} \cdots p_r^2\right]} = \frac{1}{2^{p_1^2 \cdots p_r^2 - 2P_r - p_1^2 \cdots \widehat{p_j^2} \cdots p_r^2} - 1}$$

so the lower bound becomes

$$\begin{aligned} m(r)_{[w]} &\geq \lambda_r^{n + p_1^2 \cdots p_r^2 - 1} 2^{-n} (1 - 2\lambda_r^{-p_1^2 \cdots p_r^2})^{P_r - 1} m \max_{j=1,\ldots,r} 2^{-\left[p_1^2 \cdots p_r^2 - 2P_r - 2p_1^2 \cdots \widehat{p_j^2} \cdots p_r^2\right]} \\ &= \lambda_r^{-1} (1 - 2\lambda_r^{-p_1^2 \cdots p_r^2})^{P_r - 1} m \max_{j=1,\ldots,r} \lambda_r^{n + p_1^2 \cdots p_r^2} 2^{-(n + p_1^2 \cdots p_r^2 - 2P_r - 2p_1^2 \cdots \widehat{p_j^2} \cdots p_r^2)}. \end{aligned}$$

Since $\lambda_r = 2^{\prod_{i=1}^r (1 - 1/p_i^2)}$ we have for $1 \leq i \leq r$

$$\lambda_r^{n + p_1^2 \cdots p_r^2} 2^{-\left(n + p_1^2 \cdots p_r^2 - 2P_r - 2p_1^2 \cdots \widehat{p_j^2} \cdots p_r^2\right)} = 2^{n\left(\prod_{j=1}^r (1 - 1/p_j^2) - 1\right) + P_r - p_1^2 \cdots p_r^2 + 2P_r + 2p_1^2 \cdots \widehat{p_j^2} \cdots p_r^2}. \tag{2.12}$$

Since $\prod_{j=1}^r (1 - 1/p_j^2) < 1$ and $n \leq p_1^2 \cdots p_r^2$ we have for the exponent of the last expression

$$\begin{aligned} & n\left(\prod_{j=1}^r (1 - 1/p_j^2) - 1\right) + P_r - p_1^2 \cdots p_r^2 + 2P_r + 2p_1^2 \cdots \widehat{p_j^2} \cdots p_r^2 \\ & \geq p_1^2 \cdots p_r^2 \left(\prod_{j=1}^r (1 - 1/p_j^2) - 1\right) + P_r - p_1^2 \cdots p_r^2 + 2P_r + 2p_1^2 \cdots \widehat{p_j^2} \cdots p_r^2 \\ & = 4 \prod_{j=1}^r (p_j^2 - 1) + 2p_1^2 \cdots \widehat{p_j^2} \cdots p_r^2 - 2p_1^2 \cdots p_r^2. \end{aligned} \tag{2.13}$$

I claim that this final expression is positive, which implies that the corresponding power of 2 on the right side of eq. (2.12) is greater than 1. In fact I claim that we already have $p_1^2 \cdots p_r^2 < 2 \prod_{j=1}^r (p_j^2 - 1)$; for dividing both sides by the term on the left this is equivalent to

$$2 \prod_{j=1}^r \left(1 - \frac{1}{p_j^2}\right) > 1.$$

Since the product is at least $6/\pi^2$ and $\pi^2 < 12$, this is indeed the case.

Therefore, eq. (2.12) shows that $\lambda_r^{n + p_1^2 \cdots p_r^2} 2^{-(n + p_1^2 \cdots \widehat{p_i^2} \cdots p_r^2)} \geq 1$ for $i > 1$, and consequently $m(r)_{[w]} \geq \lambda_r^{-1} (1 - 2\lambda_r^{-p_1^2 \cdots p_r^2})^{\prod_{i=1}^r (p_i^2 - 1) - 1} m$ by the previous computations. Now it is a simple calculus exercise to show that $(1 - 2^{1-x})^{x-1} \to 1$ as $x \to \infty$. Thus if we let $M = \sup_{x>1} (1 - 2^{1-x})^{x-1}$, we have $M < \infty$ and $m(r)_{[w]} \geq \lambda^{-1} M m$ for any $r \geq 1$, so $L = \lambda^{-1} M m$ satisfies the claim. □

3. THE FULL SQUAREFREE FLOW

3.1. **Topological properties.** We now turn to the full squarefree flow $\mathcal{A} = \cap_{r=1}^{\infty} \mathcal{A}(r)$. Prop. 4.4.6 of [LM] states that if $X_1 \supseteq X_2 \supseteq X_3 \supseteq \cdots$ are shift spaces whose intersection is X, then $h_{\text{top}}(X_k) \to h_{\text{top}}(X)$. Thus we have by Prop. 2.3

$$\begin{aligned} h_{\text{top}}(\mathcal{A}) &= \lim_r h_{\text{top}}(\mathcal{A}(p_1, \ldots, p_r)) \\ &= \lim_r \log 2 \prod_{p \le p_r} \left(1 - \frac{1}{p^2}\right) \\ &= \frac{6}{\pi^2} \log 2. \end{aligned}$$

More generally, for any infinite sequence of intgers $\vec{k} = (k_p) \in \prod_p \{1, \ldots, p^2 - 1\}$ (so k_p is the number of residue classes omitted modp^2) we may define $\mathcal{A}_{\vec{k}} = \cap_p \mathcal{A}_{k_p}(p)$ and $\mathcal{C}_{\vec{k}} = \cap_p \mathcal{C}_{k_p}(p)$, and then

$$\begin{aligned} h_{\text{top}}(\mathcal{C}_{\vec{k}}) &= \lim_r h_{\text{top}}(\mathcal{C}_{(k_1, \ldots, k_r)}(p_1, \ldots, p_r)) \\ &= \lim_r \log 2 \prod_{p \le p_r} \left(1 - \frac{k_p}{p^2}\right) \\ &= \log 2 \prod_p \left(1 - \frac{k_p}{p^2}\right). \end{aligned}$$

This suggests that any invariant measure of maximal entropy on $\mathcal{A}$ is concentrated on $\mathcal{A}_1$.

Lemma 3.1. *Let η be an ergodic measure on $\mathcal{A}$. Then there exists a unique sequence $\vec{k} \in \prod_p \{1, \ldots, p^2 - 1\}$ such that η is supported on $\mathcal{A}_{\vec{k}}$. In particular, any invariant (not necessarily ergodic) measure of maximal entropy on $\mathcal{A}$ is supported on $\mathcal{A}_1$.*

Proof. Although one might be tempted to reach the desired conclusion immediately from the fact that $\mathcal{A}$ is the disjoint union of the $\mathcal{A}_{\vec{k}}$ over all sequences $\vec{k}$, this is invalid since there are uncountably many such sequences, so we must argue differently. Let η be an ergodic measure on $\mathcal{A}$, and let $X = \text{supp}(\eta)$. For each $x \in X$ and each prime p let $k_p(x) = p^2 - |\pi_p(\text{supp}(x))|$. $k_p : X \to \mathbb{N}$ is measurable, and we have for any p

$$X = \coprod_{k=1}^{p^2-1} X_k(p)$$

where $X_k(p) = \{x \in X : k_p(x) = k\} = X \cap \mathcal{A}_k(p)$. Since η is ergodic with $\eta(X) = 1$, and the $X_k(p)$ are invariant (up to absolute measure zero) and mutually disjoint, there exists a unique $k_X(p) \in \{1, \ldots, p^2 - 1\}$ such that $\eta(X_{k_X(p)}) = 1$. As this is true for all p, and since a countable intersection of sets of full measure also has full measure, we have $\eta(X \cap \mathcal{A}_{\vec{k}}) = \eta\left(\cap_p X_{k_X(p)}\right) = 1$ where $\vec{k} = (k_X(p))$, which proves the first claim. The second claim follows from ergodic decomposition as in the remarks following Prop. 2.2. □

We now prove that $\mathcal{A}$ isn't sofic by bringing out the role of periodic points. By the remarks preceding Thm. 6.1.9 of [Kit], the set of periodic points in an irreducible (=topologically transitive) sofic system is dense. We will prove that $\mathcal{A}$ is irreducible but that it contains only one periodic point, so it certainly can't be sofic. Actually, since irreducibility of a subshift is equivalent to topological transitivity, and $\mathcal{A}$ is by definition the closure of the orbit of the point $(\mu^2(n))_{n \ge 1} \in \Omega^{(2)}$, it is immediate that $\mathcal{A}$ is irreducible. Nonetheless, we present a proof that uses only the definition of $\mathcal{A}$ as the space of admissible sequences.

Proposition 3.2. *$\mathcal{A}$ is irreducible.*

Proof. First, observe that if w is a finite word on $\{0,1\}$, then w appears in $\mathcal{A}$ if and only if $\pi_p(\mathrm{supp}(w))$ is a proper subset of $\mathbb{Z}/p^2\mathbb{Z}$ for every p. Indeed, the analagous claim for $\mathcal{A}(p_1,\dots,p_r)$ is immediate from the Chinese remainder theorem, and it then follows for the full squarefree flow because $\Omega^{(2)}$ is compact. To be exact, if $\pi_p(\mathrm{supp}(w))$ is proper for every prime p, then for each r we may find some $y_r \in \mathcal{A}(p_1,\dots,p_r)$ such that w appears as the first subword of y_r. Since $\Omega^{(2)}$ is compact, some subsequence of y_r converges to an element y of $\Omega^{(2)}$, which must in fact belong to $\mathcal{A}$ because $\mathcal{A}(p_1,\dots,p_{r+1}) \subset \mathcal{A}(p_1,\dots,p_r)$ for all $r \geq 1$. Since w is clearly the first subword of y, we see that $w \in \mathcal{W}(\mathcal{A})$ as desired.

As before, let A, B, A', B' be finite subsets of $\mathbb{N}$ such that $U := C_{A,B} \cap \mathcal{A}$ and $V := C_{A',B'} \cap \mathcal{A}$ are both nonempty. The key point is that since A and A' are finite, the reduction modp^2 of their union will automatically be a proper subset of $\mathbb{Z}/p^2\mathbb{Z}$ if $p^2 > |A| + |A'|$, so we only need to consider primes smaller than this fixed bound. Let $p_1 \leq p_2 \leq \cdots \leq p_r$ be the primes whose squares are less than or equal to $|A|+|A'|$, and choose a large m such that $\max A < \min(B' + mp_1^2 \cdots p_r^2)$ and $\max A' < \min(B + mp_1^2 \cdots p_r^2)$. As per the proof of Prop. 2.6, we may find some $n \in \mathbb{N}$ such that $\pi_{p_i}(A \cup (A' + n)) = \pi_{p_i}(A \cup (A' + n + mp_1^2 \cdots p_r^2))$ is proper for $i = 1, \dots, r$. On the other hand, for $p > p_r$ we have $p^2 > |A| + |A'| = |A| + |A' + n + mp_1^2 \cdots p_r^2|$. Therefore, $\pi_p(A \cup (A' + n + mp_1^2 cdotsp_r^2))$ is proper for every prime p, so

$$U \cap T^{-(n+mp_1^2\cdots p_r^2)} = C_{A\cup(A'+n+mp_1^2\cdots p_r^2),B\cup(B'+n+mp_1^2\cdots p_r^2)} \cap \mathcal{A}$$

is nonempty, as desired.

□

To study the periodic points of $\mathcal{A}$, we first examine those of the systems $\mathcal{A}(p_1,\dots,p_r)$. We begin with the case of a single prime.

Proposition 3.3. *Let p be any prime, and let x be a periodic point in $\mathcal{A}(p)$ not equal to $(0,0,0,\dots)$. Then the period of x is a multiple of p.*

Proof. Let m be the period of x. Then there exists a word $w = w_1 \cdots w_m$ such that $x = (w_1, w_2, \dots, w_m, w_1, w_2, \dots)$. Choose $a \in \{1, 2, \dots, p^2\}$ such that $a(p^2) \cap \mathrm{supp}(x) = \emptyset$, and choose any $b \in \{1, 2, \dots, m\}$ such that $w_b = 1$ (such a b exists because $x \neq (0,0,0,\dots)$). If $(m, p^2) = 1$ (which is the case if and only if m is not a multiple of p), then by the Chinese remainder theorem, we may find some $n \in \mathbb{N}$ such that $n \equiv a \bmod p^2$ and $n \equiv b \bmod m$. Therefore, there exist $\ell, \ell' \geq 0$ such that $n = a + \ell p^2 = b + \ell' m$, which implies both

$$x(n) = x(a + \ell p^2) = 0 \text{ since } a + \ell p^2 \in a(p^2)$$

and, since x has period m,

$$x(n) = x(b + \ell' m) = x(b) = w_b = 1$$

which is a contradiction.

□

This may be extended to a finite set of primes in the obvious way.

Proposition 3.4. *Let $p_1, \dots, p_r$ be distinct primes, and let x be a periodic point in $\mathcal{A}(p_1, \dots, p_r)$ not equal to $(0,0,0,\dots)$. Then the period of x is a multiple of $p_1 \cdots p_r$.*

Proof. Since $\mathcal{A}(p_1,\dots,p_r) = \cap_{i=1}^r \mathcal{A}(p_i)$, x is a periodic point of the system $\mathcal{A}(p_i)$ for $i = 1,\dots,r$, and its periods in each of these systems are all equal to its period m in $\mathcal{A}(p_1,\dots,p_r)$. Therefore, by the above proposition, m is a multiple of p_i for $i = 1,\dots,r$, so m is also a multiple of $p_1 \cdots p_r$. ☐

Prop. 3.2 and the above remarks now imply the following.

Corollary 3.5. $(0, 0, 0, \dots)$ *is the only periodic point of* $\mathcal{A}$. *In particular,* $\mathcal{A}$ *is not sofic.*

Proof. If x is a periodic point of $\mathcal{A}$, then it is also a periodic point of $\mathcal{A}(p_1,\dots,p_r)$ for all $r \geq 1$. By the above proposition, the period of x must then be a multiple of $p_1 \cdots p_r$ for all $r \geq 1$, which is absurd. ☐

To summarize, $\mathcal{A}$ is irreducible and neither sofic nor minimal since it contains the fixed point $(0, 0, 0, \dots)$. Now, it is well-known that the set of transitive points (points with dense orbit) in an irreducible system is a dense G_δ, while the set of intransitive points is either empty or dense (cf. [KS] Thm. 4.3.1). Since we know that $\mathcal{A}$ contains the intransitive point $(0, 0, 0, \dots)$, we get the following.

Corollary 3.6. *Let* $\mathcal{F}$ *be the set of intransitive points of* $\mathcal{A}$.

1. $\mathcal{F}$ *is a dense, meagre* F_σ *set.*
2. *The closure of the* T*-orbit of any point in* $\mathcal{F}\backslash\{(0, 0, 0, \dots)\}$ *is an infinite proper subshift of* $\mathcal{A}$.

One suspects that $\mathcal{F}$ carries some meaningful information relating to arithmetic progressions modulo squares of primes, but we won't pursue that possibility here.

3.2. **Approximating $\mathcal{A}$ by the finite-level systems.** Although $\mathcal{A}$ is not sofic (and doesn't fall into any class of intrinsically ergodic systems we're aware of), we are able to prove in the next section that $\mathcal{A}$ is intrinsically ergodic by showing that it is (in more than one sense) the limit of the intrinsically ergodic systems $\mathcal{A}(r)$. To begin, note that some subsequence of the measures η_r converges weak* to a measure η on $\Omega^{(2)}$, which automatically is supported on $\mathcal{A}$ and has maximal entropy by the upper semi-continuity of the entropy function. To prove that η is the only measure of maximal entropy on $\mathcal{A}$, we will exhibit it as a more refined kind of limit of the measures η_r, namely their limit under Ornstein's $\overline{d}$ metric. Good general references on the $\overline{d}$-distance are [O], [Gla] Ch. 15.7, and [Sh] Ch. I.9.

We should point out that the facts to follow only apply to invertible processes, i.e. two-sided shifts. There are extensions of Ornstein's theory to one-sided shifts (e.g. [HR]) that would apply to the systems we consider. However, we will not need these, since there is a natural entropy-preserving bijection between invariant measures on a one-sided shift and those on its invertible extension ([CT] Prop. 2.1). In particular intrinsic ergodicity for the two are equivalent, so for our applications it suffices to consider the two-sided version of $\boldsymbol{\mathcal{A}}$, i.e. in the following discussion we consider *bi*-infinite sequences of 0's and 1's with admissible support. By abuse of notation we still refer to this subshift as $\mathcal{A}$, and the associated maximal entropy measure system as $\boldsymbol{\mathcal{A}}$ (and similarly for the finite level systems $\boldsymbol{\mathcal{A}}(r)$).

We follow [Gla] §15.7. A pair $(\mathbf{X}, \alpha)$ where $\mathbf{X} = (X, S, \eta)$ is a measure-preserving dynamical system and $\alpha = \{A_i\}_{1\leq i\leq \ell}$ is a finite partition of X into disjoint measurable subsets is called a process. Associated to a process is a homomorphism $\phi_\alpha : \mathbf{X} \to (\Omega(\ell), T, (\phi_\alpha)_*\eta)$ defined by $\phi_\alpha x(n) = i \Leftrightarrow S^n x \in A_i$. We call $\rho(\mathbf{X}, \alpha) := (\phi_\alpha)_*\eta$ the symbolic representation measure of the process.

Given two processes $(\mathbf{X}_j, \alpha_j), j = 1, 2$ where both partitions have the same number of sets ℓ, define the $\bar{d}$-distance between them by

$$\bar{d}((\mathbf{X}_1, \alpha_1), (\mathbf{X}_2, \alpha_2)) = \inf\{\lambda(\alpha_1 \times X_2 \triangle X_1 \times \alpha_2) : \lambda \in J(\eta_1, \eta_2)\}$$

where $J(\eta_1, \eta_2)$ is the set of joinings of the two systems. One can show that

$$\bar{d}((\mathbf{X}_1, \alpha_1), (\mathbf{X}_2, \alpha_2)) = \bar{d}(\rho_1, \rho_2)$$

where $\rho_1 = \rho(\mathbf{X}_1, \alpha_1),\ \rho_2 = (\mathbf{X}_2, \alpha_2)$, and we note that ρ_1 and ρ_2 are both measures on $\Omega(\ell)$.

$\bar{d}$ is a complete metric on the set of T-invariant measures $M_T(\Omega)$, and the topology it induces is (much) stronger than the weak* topology ([Gla] Prop. 15.20). In order to prove that η is unique (which we accomplish in the next section), we must first show that it is ergodic in order to follow the outline of Weiss's argument for the uniqueness of the measure of maximal entropy on an irreducible sofic shift. This is supplied by the following fact.

Proposition 3.7 ([Sh] Thm. I.9.15)**.** *A $\bar{d}$-limit of ergodic measures on a shift space is ergodic.*

It is far from true that a weak* limit of ergodic measures is ergodic in general; in fact, the ergodic measures on a shift space are weak* dense in the space of invariant measures. Since each η_r is ergodic, we naturally aim to prove the following.

Theorem 3.8. *In the above notation, we have $\bar{d}(\eta_r, \eta) \to 0$ as $r \to \infty$. Therefore, η is ergodic by the above proposition.*

Actually, we will show that the sequence of processes $\boldsymbol{\mathcal{A}}(r)$ (with the time-one partition) is Cauchy under $\bar{d}$, and therefore convergent since $\bar{d}$ is complete. It follows that the entire sequence of measures η_r converges weak*, and thus its ($\bar{d}$ and weak*) limit must equal η.

To do this, we will produce for each pair of integers $r < s$ a joining ρ_{rs} of the systems $\boldsymbol{\mathcal{A}}(r)$ and $\boldsymbol{\mathcal{A}}(s)$ such that

$$\rho_{rs}(C_0^r \times C_1^s \amalg C_1^r \times C_0^s) \to 0 \text{ as } r, s \to \infty$$

where $C_0^r = \{x \in \mathcal{A}(r) : x_1 = 0\}, C_1^s = \{x \in \mathcal{A}(s) : x_1 = 1\}$ and similarly for C_1^r, C_0^s. Since $C_0^r \times C_1^s \amalg C_1^r \times C_0^s = \alpha^r \times \mathcal{A}(s) \triangle \mathcal{A}(r) \times \alpha^s$, where $\alpha^r = \{C_0^r, C_1^r\}, \alpha^s = \{C_0^s, C_1^s\}$ are the time-one partitions of $\mathcal{A}(r)$ resp. $\mathcal{A}(s)$, the definition of the $\bar{d}$ distance as an infimum over joinings implies that

$$\bar{d}(\boldsymbol{\mathcal{A}}(r), \boldsymbol{\mathcal{A}}(s)) \le \rho_{rs}(C_0^r \times C_1^s \amalg C_1^r \times C_0^s) \to 0 \text{ as } r, s \to \infty$$

which yields the desired conclusion.

Let's first sketch the idea of how to construct the joining ρ_{rs}. A sequence $x \in \Omega^{(2)}$ belongs to $\mathcal{A}(r)$ if and only if x omits at least one arithmetic progression modulo p_i^2 for $i = 1, \dots, r$, i.e. if and only if there exists $a \in \{1, 2, \dots, p_i^2\}$ such that $x_{a+\ell p_i^2} = 0$ for any $\ell \ge 0$. Outside of these arithmetic progressions, however, the entries of x may be chosen with complete freedom. Thus, suppose we begin with an r-tuple $(g_1, g_2, \dots, g_r) \in \mathbb{Z}_r := \mathbb{Z}/p_1^2\mathbb{Z} \times \cdots \times \mathbb{Z}/p_r^2\mathbb{Z}$ and *any* sequence $x' \in \Omega^2$. To define a sequence in $\mathcal{A}(r)$ from this information, we start reading the entries of x' until we reach a position congruent to one of the g_i mod p_i^2, where we insert a zero. We then continue to read x' from where we left off until we reach another position congruent to some (likely to be different) g_j mod p_j^2, where we again insert a zero, and so on. For example, in the case $r = 1$, suppose we are given the residue class 3 mod 4 and the sequence $x \in \Omega^{(2)}$ shown below. Then modifying x as just described will yield the sequence x' below:

$$\begin{aligned} x &= 01110110101101\cdots \\ x' &= 01\underline{0}110\underline{0}110\underline{0}101\underline{0}101\underline{0}\cdots \end{aligned}$$

where the underlined zeros have been inserted to force x' to omit the progression 3 mod 4.

We can formalize this as follows. Given any $g = (g_1, \ldots, g_r) \in \mathbb{Z}_r$, define a map $\alpha_g : \mathbb{N} \to \mathbb{N}$ by $\alpha_g(n) = |\{m \leq n : m \equiv g_i \bmod p_i^2 \text{ for some } i \leq r\}|$. Then define $\phi_r : \mathbb{Z}_r \times \Omega^{(2)} \to \Omega^{(2)}$ by

$$\phi_r(g, x)_n = \begin{cases} 0 & \text{if } n \equiv g_i \bmod p_i^2 \text{ for some } i \leq r \\ x_{n-\alpha_g(n)} & \text{else} \end{cases}$$

It is straightforward to verify that ϕ_r is continuous and intertwines the $\mathbb{Z}$-actions on the two systems when we consider translation by $(-1, -1, \ldots, -1)$ on $\mathbb{Z}_r$. Moreover, the image of ϕ_r equals $\mathcal{A}(r)$ by construction. Therefore ϕ_r is a topological factor map from $\mathbb{Z}_r \times \Omega^{(2)}$ onto $\mathcal{A}(r)$.

The following proposition reflects the fact that the entries of a member of $\mathcal{A}(r)$ may be chosen freely once it's omitted at least one arithmetic progression modp_i^2 for each i.

Proposition 3.9. *Define the measure* $\nu_r := m_r \times B(\frac{1}{2}, \frac{1}{2})$ *on* $\mathbb{Z}_r \times \Omega^{(2)}$ *where* m_r *is the counting measure and* $B(\frac{1}{2}, \frac{1}{2})$ *is the maximum entropy Bernoulli measure on* $\Omega^{(2)}$*. Then* $(\phi_r)_*\nu_r = \eta_r$*, the measure of maximal entropy on* $\mathcal{A}(r)$*.*

Proof. Let $\omega = (\phi_r)_*\nu_r$; this is an invariant measure on $\mathcal{A}(r)$. We will show that the measure entropy $h_\omega(\mathcal{A}(r)) = h_{\text{top}}(\mathcal{A}(r))$, which implies $\omega = \eta_r$ since η_r is the only measure of maximal entropy on $\mathcal{A}(r)$ by prop. 2.6.

To this end, let $n \geq 1$ and consider a word $w \in \mathcal{W}(\mathcal{A}(r))$ of length n. Let $A_w = \mathcal{A}(r)_w$ be the associated cylinder set and let $B_w = \phi_r^{-1}(A_w)$, so that $\omega(A_w) = \nu_r(B_w)$. Define

$$\begin{aligned} I_0 &= \{m \leq n : w_m = 0\} \\ I_1 &= \{m \leq n : w_m = 1\}. \end{aligned}$$

By definition, we have $(g, x) \in B_w$ if and only if $\phi_r(g, x)_m = w_m$ for $m = 1, \ldots, n$. Thus if $m \in I_0$, we must have $x_{m-\alpha_g(m)} = 0$ or $m \equiv g_i \bmod p_i^2$ for some $i \leq r$, while if $m \in I_1$ we must have $x_{m-\alpha_g(m)} = 1$ and $m \not\equiv g_i \bmod p_i^2$ for any $i \leq r$. Defining $\delta_g(m) = 1$ if and only if $m \equiv g_i \bmod p_i^2$ for some $i \leq r$ and $\delta_g(m) = 0$ else, it follows that we have

$$\begin{aligned} B_w = \Big\{ &(g, x) \in \mathbb{Z}_r \times \Omega^{(2)} : x_{m-\alpha_g(m)} = 0 \ \forall m \in I_0 \text{ with } \delta_g(m) = 0, \\ &x_{m-\alpha_g(m)} = 1 \ \forall m \in I_1, \text{ and } g_i \not\equiv m \bmod p_i^2 \text{ for any } m \in I_1, i \leq r \Big\}. \end{aligned}$$

Assume without loss of generality that n, the length of w, has form $n = kp_1^2 \cdots p_r^2$. We will show that for any $g \in \mathbb{Z}_s$ we have

$$\big\{m - \alpha_g(m) : m \in \big[1, kp_1^2 \cdots p_r^2\big] \text{ with } \delta_g(m) = 0\big\} = \big[1, k(p_1^2 - 1) \cdots (p_r^2 - 1)\big] \tag{3.1}$$

where $[a, b]$ denotes the set of integers between a and b. For this, define a map $\beta_g : \mathbb{N} \to \mathbb{N}$ by $\beta_g(m) = \min\{\ell > m : \delta_g(\ell) = 0\}$. I claim that $\beta_g(m) - \alpha_g(\beta_g(m)) = m + 1 - \alpha_g(m)$ for any m. Indeed, the number of ℓ strictly between m and $\beta_g(m)$ with $\delta_g(\ell) = 1$ is by definition $\alpha_g(\beta_g(m)) - \alpha_g(m)$. Since $\beta_g(m)$ by construction equals $m + 1$ plus the number of such ℓ, we see that $\beta_g(m) = m + 1 + \alpha_g(\beta_g(m)) - \alpha_g(m)$ as claimed.

Define numbers β_k for $k = 1, \dots, n - \alpha_g(n)$ by $\beta_1 = 1$ if $\delta_g(1) = 0$ and $\beta_1 = \beta_g(1)$ otherwise, then set $\beta_k = \beta_g(\beta_{k-1})$ for $k > 1$. Then it is clear that $\delta_g(m) = 0$ if and only if there exists some k such that $m = \beta_k$. Therefore

$$\begin{aligned}&\{m - \alpha_g(m) : m \in \left[1, kp_1^2 \cdots p_r^2\right] \text{ with } \delta_g(m) = 0\}\\ &= \left\{\beta_1 - \alpha_g(\beta_1), \beta_2 - \alpha_g(\beta_2), \dots, \beta_{n-\alpha_g(n)} - \alpha_g(\beta_{n-\alpha_g(n)})\right\}.\end{aligned}$$

By the above remarks, we have $\beta_{k+1} = \beta_k + 1 + \alpha_g(\beta_{k+1}) - \alpha_g(\beta_k)$, which is equivalent to $(\beta_{k+1} - \alpha_g(\beta_{k+1})) - (\beta_k - \alpha_g(\beta_k)) = 1$ for all k; therefore

$$\left\{\beta_1 - \alpha_g(\beta_1), \beta_2 - \alpha_g(\beta_2), \dots, \beta_{n-\alpha_g(n)} - \alpha_g(\beta_{n-\alpha_g(n)})\right\} = [1, n - \alpha_g(n)].$$

Since $n = kp_1^2 \cdots p_r^2$ and $n - \alpha_g(n)$ equals the number of $m \le n$ with $\delta_g(m) = 0$, we see by inclusion-exclusion and the Chinese remainder theorem that $n - \alpha_g(n) = k(p_1^2 - 1) \cdots (p_r^2 - 1)$, which implies the claim.

By the above decomposition of B_w we have

$$B_w = \coprod_{g \in G_1} \{g\} \times C_{\alpha_g}$$

where $G_1 = \{g \in \mathbb{Z}_s : g_i \not\equiv m \bmod p_i^2 \text{ for any } m \in I_1, i \le r\}$ and $C_{\alpha_g} \subset \Omega^{(2)}$ is the subset $\{x \in \Omega^{(2)} : x_{m-\alpha_g(m)} = 0 \ \forall m \in I_0 \text{ with } \delta_g(m) = 0 \text{ and } x_{m-\alpha_g(m)} = 1 \ \forall m \in I_1\}$. Therefore, by eq. (3.1) and the definition of the Bernoulli measure we have

$$\begin{aligned}\nu_r(B_w) &= \frac{1}{p_1^2 \cdots p_r^2} \sum_{g \in G_1} 2^{-k \prod_{i=1}^r (p_i^2 - 1)}\\ &= \frac{1}{p_1^2 \cdots p_r^2} 2^{-k \prod_{i=1}^r (p_i^2 - 1)} |G_1|\\ &= \frac{1}{p_1^2 \cdots p_r^2} 2^{-k \prod_{i=1}^r (p_i^2 - 1)} \prod_{i=1}^r (p_i^2 - u(w,i))\\ &= 2^{-k \prod_{i=1}^r (p_i^2 - 1)} \prod_{i=1}^r (1 - u(w,i)/p_i^2)\end{aligned}$$

where $u(w,i)$ is the number of distinct residue classes modulo p_i^2 defined by I_1. Letting α be the time one partition on $\mathcal{A}(r)$ and $n = kp_1^2 \cdots p_r^2$ as above, it follows that

$$\begin{aligned}H_\omega(\alpha_0^{n-1}) &= - \sum_{\substack{w \in \mathcal{W}(\mathcal{A}(r))\\ |w| = n}} \nu_r(B_w) \log \nu_r(B_w)\\ &= \sum_{\substack{w \in \mathcal{W}(\mathcal{A}(r))\\ |w| = n}} 2^{-k \prod_{i=1}^r (p_i^2 - 1)} \prod_{i=1}^r (1 - u(w,i)/p_i^2) \left[k \prod_{i=1}^r (p_i^2 - 1) \log 2 - \sum_{i=1}^r \log(1 - u(w,i)/p_i^2)\right]\end{aligned}$$

We will focus on the case when $u(w,i) = p_i^2 - 1$ for $i = 1, \dots, r$. Indeed, we see from the above and the fact that $\sum_{i=1}^r \log(1 - u(w,i)/p_i^2) < 0$ that

$$\begin{aligned}
H_\omega(\alpha_0^{n-1}) &\geq 2^{-k\prod_{i=1}^r (p_i^2-1)} \sum_{\substack{w\in\mathcal{W}(\mathcal{A}(r))\\ |w|=n \\ u(w,i)=p_i^2-1 \text{ for } i=1,\dots,r}} \prod_{i=1}^r (1-(p_i^2-1)/p_i^2)\left[k\prod_{i=1}^r (p_i^2-1)\log 2\right] \\
&= 2^{-k\prod_{i=1}^r (p_i^2-1)} \sum_{\substack{w\in\mathcal{W}(\mathcal{A}(r))\\ |w|=n \\ u(w,i)=p_i^2-1 \text{ for } i=1,\dots,r}} \prod_{i=1}^r 1/p_i^2\left[k\prod_{i=1}^r (p_i^2-1)\log 2\right] \\
&= k2^{-k\prod_{i=1}^r (p_i^2-1)} \sum_{\substack{w\in\mathcal{W}(\mathcal{A}(r))\\ |w|=n \\ u(w,i)=p_i^2-1 \text{ for } i=1,\dots,r}} \log 2\prod_{i=1}^r \left(1-\frac{1}{p_i^2}\right) \\
&= k2^{-k\prod_{i=1}^r (p_i^2-1)} \times \left|\left\{w\in\mathcal{W}(\mathcal{A}(r)) : |w| = kp_1^2\cdots p_r^2 \text{ and } u(w,i)=p_i^2-1 \text{ for } i=1,\dots,r\right\}\right| \\
&\qquad \times \log 2\prod_{i=1}^r \left(1-\frac{1}{p_i^2}\right).
\end{aligned}$$

Since by definition $H_\omega(\alpha_0^{kp_1^2\cdots p_r^2-1})/(kp_1^2\cdots p_r^2) \to h_\omega(\mathcal{A}(r))$ as $k\to\infty$, and since $h_\omega(\mathcal{A}(r)) \leq h_{\text{top}}(\mathcal{A}(r))$ by the variational principle, we see by recognizing $h_{top}(\mathcal{A}(r))$ as the last factor in the product above that we'll be done if we can show

$$\begin{aligned}
\lim_{k\to\infty} 2^{-k\prod_{i=1}^r (p_i^2-1)} \cdot \left|\left\{w\in\mathcal{W}(\mathcal{A}(r)) : |w| = kp_1^2\cdots p_r^2 \text{ and } u(w,i)=p_i^2-1 \text{ for } i=1,\dots,r\right\}\right| \\
\geq p_1^2\cdots p_r^2. \qquad (3.2)
\end{aligned}$$

First consider the case $r=1$, so that we are considering just one prime p. Let

$$W^1(k) = \left\{w\in\mathcal{W}(\mathcal{A}(p)) : |w| = kp^2 \text{ and } u(w) = p^2-1\right\},$$

that is to say, $W^1(k)$ consists of those words of length kp^2 whose support omits exactly one residue class modp^2. We have

$$W^1(k) = \coprod_{g\in\mathbb{Z}/p^2\mathbb{Z}} W_g^1(k)$$

where $W_g^1(k)$ is the set of length kp^2 words that omit only g mod p^2 from their support.

Now, for each $g\in\mathbb{Z}/p^2\mathbb{Z}$ let

$$W_g(k) = \{w\in\mathcal{W}(\mathcal{A}(p)) : |w| = kp^2 \text{ and } g\notin\pi_p(\text{supp}(w))\}.$$

That is, $W_g(k)$ consists of all length kp^2 words that omit g mod p^2 (but may omit other residue classes as well). Then we have $W_g(k) = W_g^1(k)\coprod V_g(k)$, where $W_g^1(k)$ is as above and $V_g(k)$ is the set of length kp^2 words that omit g and at least one other progression modp^2.

Observe that $|W_g(k)| = 2^{kp^2-k} = 2^{k(p^2-1)}$, since $W_g(k)$ is defined solely by the conditions that w have length kp^2 and have zeros in the positions $\widetilde{g}+\ell p^2$ for $\ell = 0,1,\dots,k-1$, where $\widetilde{g}$ is the smallest integer representative of $g\in\mathbb{Z}/p^2\mathbb{Z}$. On the other hand, we have

$$V_g(k) = \bigcup_{\substack{h\in\mathbb{Z}/p^2\mathbb{Z}\\ h\neq g}} \{w\in W_g(k) : h\notin\pi_p(\text{supp}(w))\}.$$

For any $h \in \mathbb{Z}/p^2\mathbb{Z}, h \neq g$, the condition that w belong to the set $\{w \in W_g(k) : h \notin \pi_p(\mathrm{supp}(w))\}$ determines $2k$ entries of w; namely, that w must have zeros in the positions $\widetilde{g} + \ell p^2$ and $\widetilde{h} + \ell p^2$ for $\ell = 0, 1, \ldots, k-1$. The other entries of w may be chosen with complete freedom; therefore $|\{w \in W_g(k) : h \notin \pi_p(\mathrm{supp}(w))\}| = 2^{kp^2-2k} = 2^{k(p^2-2)}$. As this is independent of the choice of h we get the estimate

$$|V_g(k)| \leq (p^2-1)2^{k(p^2-2)}.$$

Hence, the decomposition $W_g(k) = W_g^1(k) \coprod V_g(k)$ for any $g \in \mathbb{Z}/p^2\mathbb{Z}$ yields the lower bound

$$|W_g^1(k)| = |W_g(k)| - |V_g(k)| \geq 2^{k(p^2-1)} - (p^2-1)2^{k(p^2-2)}$$

and it follows that for any g we have

$$\lim_{k\to\infty} 2^{-k(p^2-1)}|W_g^1(k)| \geq 1.$$

Since this holds for any g, we find

$$\lim_{k\to\infty} 2^{-k(p^2-1)}|W^1(k)| = \sum_{g\in\mathbb{Z}/p^2\mathbb{Z}} \lim_{k\to\infty} 2^{-k(p^2-1)}|W_g^1(k)| \geq p^2$$

which proves the inequality (3.2) in this case.

In fact, we can deduce the inequality (3.2) for any $r \geq 1$ from this using induction on r and the Chinese remainder theorem. Indeed, let

$$W_r^1(k) = \left\{w \in \mathcal{W}(\mathcal{A}(r)) : |w| = kp_1^2 \cdots p_r^2 \text{ and } u(w,i) = p_i^2 - 1 \text{ for } i = 1, \ldots, r\right\}.$$

Our aim is to show that $\lim_{k\to\infty} 2^{-k\prod_{i=1}^r (p_i^2-1)}|W_r^1(k)| \geq p_1^2 \cdots p_r^2$.

A word w of length $kp_1^2 \cdots p_r^2$ belongs to $W_r^1(k)$ if and only if w belongs to $W_{r-1}^1(kp_r^2)$ and the support of w omits exactly one residue class $\bmod\, p_r^2$. The latter condition forces w to have zeros in the $kp_1^2 \cdots p_{r-1}^2$ positions $\tilde{g} + \ell p_r^2$ for $\ell = 0, 1, \ldots, kp_1^2 \cdots p_{r-1}^2 - 1$, where $\pi_r(\tilde{g}) \in \mathbb{Z}/p_r^2\mathbb{Z}$ is the unique class omitted by $w \bmod p_r^2$. By the Chinese remainder theorem, this implies that

$$|W_r^1(k)| = p_r^2 2^{-k\prod_{i=1}^{r-1}(p_i^2-1)}|W_{r-1}^1(kp_r^2)| + O(1).$$

Consequently

$$\begin{aligned} 2^{-k\prod_{i=1}^r (p_i^2-1)}|W_r^1(k)| &= 2^{-kp_r^2\prod_{i=1}^{r-1}(p_i^2-1) + k\prod_{i=1}^{r-1}(p_i^2-1)}|W_r^1(k)| \\ &= p_r^2 2^{-kp_r^2\prod_{i=1}^{r-1}(p_i^2-1)}|W_{r-1}^1(kp_r^2)| + o(1) \text{ as } k \to \infty. \end{aligned}$$

The claim now follows by induction on r, the case $r = 1$ having been shown above. Indeed, assuming the lower bound (3.2) holds for $r-1$, we have

$$\begin{aligned} \lim_{k\to\infty} 2^{-k\prod_{i=1}^r (p_i^2-1)} \cdot \left|W_r^1(k)\right| &= \lim_{k\to\infty} p_r^2 2^{-kp_r^2\prod_{i=1}^{r-1}(p_i^2-1)} \left|W_{r-1}^1(kp_r^2)\right| \\ &\geq p_r^2 \cdot p_1^2 \cdots p_{r-1}^2. \end{aligned}$$

Now it follows from the computations preceding the inequality (3.2) that $h_\omega(\mathcal{A}(r)) = h_{\mathrm{top}}(\mathcal{A}(r))$, which shows that $\omega = \eta_r$, as desired. □

Now suppose $s > r$, so we have maps $\phi_s : \mathbb{Z}_s \times \Omega^{(2)} \to \mathcal{A}(s)$ and $\phi_r : \mathbb{Z}_r \times \Omega^{(2)} \to \mathcal{A}(r)$. Since $s > r$, $\mathbb{Z}_r$ is naturally a factor of $\mathbb{Z}_s$ under the map which ignores the last $s - r$ coordinates. Therefore, we may pull back ϕ_r to a map $\widetilde{\phi}_r : \mathbb{Z}_s \times \Omega^{(2)} \to \mathcal{A}(r)$. This remains a topological factor map, and we clearly have $(\widetilde{\phi}_r)_*\nu_s = \eta_r$.

Denote by $\mathbf{X}_s$ the measure-preserving system $(\mathbb{Z}_s \times \Omega^{(2)}, \nu_s)$ with the action being the product of translation by $(-1, -1, \ldots, -1)$ on $\mathbb{Z}_s$ with the left shift on $\Omega^{(2)}$. Then the above constructions imply that $\mathbf{X}_s$ is an extension of both $\boldsymbol{\mathcal{A}}(r)$ and $\boldsymbol{\mathcal{A}}(s)$ under the maps $\widetilde{\phi}_r$ resp. ϕ_s. Thus, we have the joining

$$\rho_{rs} := (\widetilde{\phi}_r \times \phi_s)_* \nu_s$$

of $\boldsymbol{\mathcal{A}}(r)$ and $\boldsymbol{\mathcal{A}}(s)$.

We proceed to estimate $\rho_{rs}(C_0^r \times C_1^s \amalg C_1^r \times C_0^s) = \rho_{rs}(C_0^r \times C_1^s) + \rho_{rs}(C_1^r \times C_0^s)$. We have by definition

$$\rho_{rs}(C_0^r \times C_1^s) = \nu_s((\widetilde{\phi}_r \times \phi_s)^{-1}(C_0^r \times C_1^s)).$$

Suppose $(g, x) \in (\widetilde{\phi}_r \times \phi_s)^{-1}(C_0^r \times C_1^s)$. To have $\widetilde{\phi}_r(g, x) \in C_0^r$ we must have $x_1 = 0$ or $g_i \equiv 1 \bmod p_i^2$ for some $i \leq r$, while to have $\phi_s(g, x) \in C_1^s$ we must have $x_1 = 1$ and $g_i \not\equiv 1 \bmod p_i^2$ for $i = 1, \ldots, s$. These conditions are clearly incompatible; therefore $(\widetilde{\phi}_r \times \phi_s)^{-1}(C_0^r \times C_1^s) = \emptyset$ and $\rho_{rs}(C_0^r \times C_1^s) = 0$.

We now turn to

$$\rho_{rs}(C_1^r \times C_0^s) = \nu_s((\widetilde{\phi}_r \times \phi_s)^{-1}(C_1^r \times C_0^s)).$$

Suppose $(g, x) \in (\widetilde{\phi}_r \times \phi_s)^{-1}(C_1^r \times C_0^s)$. To have $\widetilde{\phi}_r(g, x) \in C_1^r$ we must have $x_1 = 1$ and $g_i \not\equiv 1 \bmod p_i^2$ for $i = 1, \ldots, r$ (but it is important to note that we *can* have $g_i \equiv 1 \bmod p_i^2$ for $i > r$), while to have $\phi_s(g, x) \in C_0^s$ we must have $x_1 = 0$ or $g_i \equiv 1 \bmod p_i^2$ for some $i \leq s$. The only way these conditions can be met simultaneously is if $x_1 = 1$ and $g_i \equiv 1 \bmod p_i^2$ for some $i > r$. Therefore

$$(\widetilde{\phi}_r \times \phi_s)^{-1}(C_1^r \times C_0^s) = G_{(r,s]} \times C_1$$

where $C_1 = \{x \in \Omega^{(2)} : x_1 = 1\}$ and

$$G_{(r,s]} = \left\{ g \in \mathbb{Z}_s : g_i \equiv 1 \bmod p_i^2 \text{ for some } r < i \leq s \right\}.$$

We have

$$G_{(r,s]} = \bigcup_{i=r+1}^{s} \left\{ g \in \mathbb{Z}_s : g_i \equiv 1 \bmod p_i^2 \right\}$$

and therefore

$$|G_{(r,s]}| \leq p_1^2 \cdots p_r^2 \sum_{i=r+1}^{s} p_{r+1}^2 \cdots \widehat{p_i^2} \cdots p_s^2.$$

Since $B(\frac{1}{2}, \frac{1}{2})(C_1) = 1/2$, we finally obtain

$$\begin{aligned}
\rho_{rs}(C_1^r \times C_0^s) &= \nu_s((\widetilde{\phi}_r \times \phi_s)^{-1}(C_1^r \times C_0^s)) \\
&= \frac{1}{2p_1^2 \cdots p_s^2} \left| G_{(r,s]} \right| \\
&\leq \frac{1}{2p_{r+1}^2 \cdots p_s^2} \sum_{i=r+1}^{s} p_{r+1}^2 \cdots \widehat{p_i^2} \cdots p_s^2 \\
&= \frac{1}{2} \left(\frac{1}{p_{r+1}^2} + \frac{1}{p_{r+2}^2} + \cdots + \frac{1}{p_s^2} \right)
\end{aligned}$$

and this converges to 0 as $r, s \to \infty$ because the sum of the reciprocals of the squares converges. This proves thm. 3.8.

3.3. **Intrinsic ergodicity.** We now come to the proof of our main theorem.

Theorem 3.10. *η is the only measure of maximal entropy on $\mathcal{A}$.*

Our proof of this theorem relies on the following lemma of Weiss. Recall that the intrinsic ergodicity of $\mathcal{A}$ is equivalent to that of its two-sided extension, so by abuse of notation we may consider the inverse of the shift acting on $\mathcal{A}$. With this understood, for a partition α and $n \geq 1$ we use the notation $\alpha_{-n}^{n} = \vee_{j=-n}^{n} T^{-j}\alpha$.

Lemma 3.11 ([Wei] Lemma 2)**.** *Let Y be a closed invariant subset of $\Omega^{(2)}$ with topological entropy $\log\lambda$, and let α be the time one partition of Y. Suppose η is an ergodic measure on Y such that $h_{\eta}(Y) = h_{top}(Y) = \log\lambda$, and suppose there is a constant $c > 0$ such that for any atom $A \in \alpha_{-n}^{n}$ we have $\eta(A) \geq c^{-1}\lambda^{-2n-1}$. Then Y is intrinsically ergodic.*

Thus we need to show that η satisfies the conditions of this lemma; since we proved in the last section that η is ergodic, we need only prove the existence of a constant $c > 0$ satisfying the stated lower bound. This bound is often referred to as the "lower Gibbs property" for the measure η, cf. [CT] §5.3 (I thank Dan Thompson for this terminology), and we see that it is powerful enough to imply the uniqueness of an ergodic measure of maximal entropy.

Recall the definition of the finite-level measures η_r (eq. (2.2)): given a word $w \in \mathcal{W}(\mathcal{A}(r))$ of length n we set

$$\eta_r(\mathcal{A}(r)_w) = \lambda_r^{-n} m(r)_{[w]}$$

where $m(r)$ is the strictly positive right Perron-Frobenius eigenvector for the matrix $F(r)$ described in sec. 2.3, normalized so that $m(r)_{[0]} + m(r)_{[1]} = \lambda_r$. The strict positivity of $m(r)$ then implies that the quantity $\lambda_r^{|w|} \cdot \eta_r(\mathcal{A}(r)_w)$ is uniformly bounded below as w varies over the words in $\mathcal{A}(r)$. Thus η_r satisfies the lower Gibbs bound, verifying the fact that η_r is the only measure of maximal entropy on $\mathcal{A}(r)$ (this is precisely why an irreducible sofic system always carries a unique measure of maximal entropy).

In our specific setting, we have uniform control over the lower Gibbs bound for $\mathcal{A}(r)$ as $r \to \infty$; namely, by prop. 2.8 we know there exists a constant $L > 0$ such that for any $r \geq 1$ and any word $w \in \mathcal{W}(\mathcal{A}(r))$ of length $2n+1 \leq p_1^2 \cdots p_r^2$, we have

$$\lambda_r^{2n+1} \cdot \eta_r(\mathcal{A}(r)_w) \geq L. \tag{3.3}$$

Now, we showed in the last section that η is the $\overline{d}$-limit of the measures η_r, so all the more so is their limit in the weak* topology. Consequently, if w is a word in $\mathcal{A}$ of length $2n+1$, we clearly have $\mathcal{A}_w = \cap_{r=1}^{\infty}\mathcal{A}(r)_w$ and, by general measure theory, $\eta(\mathcal{A}_w) = \lim_{r\to\infty}\eta_r(\mathcal{A}_w) = \lim_{r\to\infty}\eta_r(\mathcal{A}(r)_w)$. Therefore eq. (3.3) yields

$$\lambda^{2n+1} \cdot \eta(\mathcal{A}_w) = \lim_{r\to\infty} \lambda_r^{2n+1} \cdot \eta_r(\mathcal{A}(r)_w) \geq L$$

since w is fixed and we have $2n+1 \leq p_1^2 \cdots p_r^2$ for all sufficiently large r.

Therefore η has the lower Gibbs property. Since it is ergodic by 3.8, Weiss's lemma then implies that η is the only measure of maximal entropy on $\mathcal{A}$. This completes the proof of 3.10.

3.4. **Structure of the maximal entropy system.** For each $r \geq 1$, let $\boldsymbol{\mathcal{A}}(r)$ be the measure-preserving dynamical system $(\mathcal{A}(r), T, \eta_r)$ and let $\boldsymbol{\mathcal{A}} = (\mathcal{A}, T, \eta)$ where η is the unique measure of maximal entropy on $\mathcal{A}$. By the main result of [T], the class of processes of form $(\mathcal{Z} \times \mathcal{B}, S \times T, m \times \beta)$ where $(\mathcal{Z}, S, m)$ is ergodic of entropy zero and $(\mathcal{B}, T, \beta)$ is a Bernoulli shift is closed under taking factors. We refer to such processes as entropy zero × Bernoulli for obvious reasons. Such a system has $\mathcal{Z}$ as its Pinsker factor. In our setting, $\mathbb{Z}_r$ (equipped with translation by $(-1, -1, -1, \dots)$) is naturally a factor of $\boldsymbol{\mathcal{A}}(r)$ under the map that sends a sequence $x \in \mathcal{A}_1(r)$ to the unique residue class $\bmod p_1^2 \cdots p_r^2$ omitted by its support (recall that $\eta_r(\mathcal{A}_1(r)) = 1$). It follows that $\mathbb{Z}_r$ is a factor of the Pinsker factor of $\boldsymbol{\mathcal{A}}(r)$.

By prop. 3.9, $\boldsymbol{\mathcal{A}}(r)$ is a factor of $\mathbb{Z}_r \times \mathcal{B}$, where $\mathcal{B}$ is the Bernoulli system of entropy $\log 2$. Moreover, by the fundamental work of Ornstein, a Bernoulli shift is classified up to isomorphism by its entropy [O]. Combining this with Thouvenot's theorem and the observations above yields the following.

Lemma 3.12. *The map ϕ_r yields an isomorphism $\boldsymbol{\mathcal{A}}(r) = \mathbb{Z}_r \times \mathcal{B}_r$, where $\mathcal{B}_r$ is the Bernoulli system of entropy $\log 2 \prod_{i=1}^{r} \left(1 - 1/p_i^2\right)$. In particular, $\mathbb{Z}_r$ is the Pinsker factor of $\boldsymbol{\mathcal{A}}(r)$.*

By thm. 3.8, the processes $\boldsymbol{\mathcal{A}}(r)$ (with their time zero partitions) converge under $\bar{d}$ to the process $\boldsymbol{\mathcal{A}}$. By the main result of [ShT], the class of entropy zero × Bernoulli processes is closed under taking $\bar{d}$-limits. In our case, the entropy zero components of the systems $\boldsymbol{\mathcal{A}}(r)$ have discrete spectrum, which must increase to the spectrum of the entropy zero component of the $\bar{d}$-limit (cf. [RS]). This finally yields the following description of the maximal entropy measure system for the squarefree flow.

Theorem 3.13. *Let $\mathcal{K}$ be the Kronecker system $\left(\prod_p \mathbb{Z}/p^2\mathbb{Z}, T_{(-1,-1,-1,\cdots)}, \nu_{Haar}\right)$. Then we have an isomorphism $\boldsymbol{\mathcal{A}} = \mathcal{K} \times \mathcal{B}_{sq}$, where $\mathcal{B}_{sq}$ is the Bernoulli system of entropy $(6/\pi^2)\log 2$. In particular, $\mathcal{K}$ is the Pinsker factor of $\boldsymbol{\mathcal{A}}$.*

3.5. **Conclusion.** To summarize, we have shown that there is a unique measure of maximal entropy on the squarefree flow and described the structure of the associated maximal entropy measure-preserving system. One may naturally wonder whether these results could be obtained in a more hands-off way by proving that the squarefree flow belongs to a class of systems that are known to be intrinsically ergodic. It's easy to see that this won't be one of the common classes (sofic shifts, β-shifts, S-gap shifts, etc.), one reason being that $\mathcal{A}$ has only one periodic point. I am unaware of any classes very different from these for which intrinsic ergodicity is known. Alternately, given its arithmetic significance, the squarefree flow may be the first example of a new class of intrinsically ergodic systems that deserve to have their properties formally extracted and defined.

REFERENCES

[AShS] R. Adler, P. Shields, and M. Smorodinsky, *Irreducible Markov Shifts*, Ann. Math. Stat. **43** no. 3, 1027–1029 (1972).

[Bro] J. Brown, *Inverse limits, entropy, and weak isomorphism for discrete dynamical systems*, Trans. Amer. Math. Soc. **164**, 55 – 66 (1972).

[CoP] E. Coven and M. Paul, *Endomorphisms of irreducible subshifts of finite type*, Math. Syst. Theory **8** no. 2, 167 –175 (1974).

[CS] F. Cellarosi and Y. Sinai, *Ergodic properties of square-free numbers*, preprint, http://arxiv.org/abs/1112.4691 (2011).

[CT] V. Climenhaga and D. Thompson, *Intrinsic ergodicity beyond specification*, preprint, http://arxiv.org/abs/1011.2780 (2011).

[E] M. Ellis, *On Kamae's conjecture concerning the $\bar{d}$-distance between two-state Markov processes*, Ann. Prob. **8** no. 2, 372–376 (1980).

[Ei] M. Einsiedler et. al., *The distribution of closed geodesics on the modular surface and Duke's theorem*, available at http://math.stanford.edu/˜akshay/research/elmv2.pdf (2011).

[Fis] R. Fischer, *Sofic systems and graphs*, Monathshefte für Mathematik, **80**, 179–186 (1975).

[Gla] E. Glasner, *Ergodic theory via joinings*, Amer. Math. Soc. Mathematical Surveys and Monographs **101** (2003).

[GTW] E. Glasner, J.P. Thouvenot, B. Weiss, *Entropy theory without a past*, Erg. Theory Dyn. Syst. **20**, 1355–1370 (2000).

[Gur] B.M. Gurevic, *Stationary random sequences of maximal entropy*, "Multicomponent Random Systems", 327 – 380, ed. Dobrushin and Sinai, Adv. in Prob. and Related Topics **6**, Marcel Dekker Inc. (1980)

[Hay] N. Haydn, *Phase transitions in one-dimensional subshifts*, preprint, http://arxiv.org/abs/1111.6227 (2011).

[HR] C. Hoffmann and D. Rudolph, *Uniform endomorphisms which are isomorphic to a Bernoulli shift*, Ann. Math. **156** no. 1, 79–101 (2002).

[IK] H. Iwaniec and E. Kowalski, *Analytic number theory*, AMS Colloquium Publications **53** (2004).

[Kit] B. Kitchens, *Symbolic dynamics: One-sided, two-sided, and countable state Markov shifts*, Springer-Verlag (1998).

[KS] S. Kolyada and L. Snoha, *Some aspects of topological transitivity - a survey*, Grazer Math. Ber., **334**, 3–35 (1997).

[KiS] B. Kitchens and S. Tuncel, *Semi-groups and graphs for sofic systems*, Israel J. Math. **53** no. 2 (1986).

[LM] D. Lind and B. Marcus, *An introduction to symbolic dynamics and coding*, Cambridge University Press (1995).

[O] D. Ornstein, *Ergodic theory, randomness, and dynamical systems*, Yale Mathematical Monographs, no. 5 (1974).

[RS] D. Rudolph and G. Schwarz, *The limits in $\bar{d}$ of multi-step Markov chains*, Israel J. Math. **28**, 103–109 (1977).

[Sar] P. Sarnak, *Three lectures on the Möbius function, randomness, and dynamics*, online notes, http://publications.ias.edu/sarnak/paper/512 (2011).

[Sh] P. Shields, *The ergodic theory of discrete sample paths*, AMS Graduate Studies in Math **13** (1996).

[ShT] P. Shields and J.P. Thouvenot, *Entropy zero × Bernoulli processes are closed in the $\bar{d}$-metric*, Ann. Probability, **3** no. 4, 732–736 (1975).

[T] P. Thouvenot, *Une classe de systemes pour lesquels la conjecture de Pinsker est vraie*, Israel J. Math. **21**, 208–214 (1975).

[Tho] K. Thomsen, *On the structure of a sofic shift space*, Trans. Amer. Math. Soc. **356** no. 9, 3557–3619 (2004).

[Tsa] K.M. Tsang, *The distribution of r-tuples of squarefree numbers*, Mathematika **32**, 265-275 (1985).

[Wei] B. Weiss, *Subshifts of finite type and sofic systems*, Monathshefte für Mathematik, **77**, 462–474 (1972).

Department of Mathematics, Princeton University
E-mail address: `rpeckner@math.princeton.edu`